

表 4477 腹膜炎罹患回数(回/年) 一日透析液使用量(L/日)別(腹膜透析単独患者)

一日透析液使用量	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2未満	88	8									96		10	106
(%)	(91.7)	(8.3)									(100.0)			
2～	313	33	4	2	1		1				354	3	14	371
(%)	(88.4)	(9.3)	(1.1)	(0.6)	(0.3)		(0.3)				(100.0)			
4～	837	86	11	4	2				1		941		45	986
(%)	(88.9)	(9.1)	(1.2)	(0.4)	(0.2)				(0.1)		(100.0)			
6～	1,229	156	30	7		2					1,424	2	67	1,493
(%)	(86.3)	(11.0)	(2.1)	(0.5)		(0.1)					(100.0)			
8～	852	94	19	4	2	1					972	15	54	1,041
(%)	(87.7)	(9.7)	(2.0)	(0.4)	(0.2)	(0.1)					(100.0)			
10～	223	36	4	3	1						267	1	5	273
(%)	(83.5)	(13.5)	(1.5)	(1.1)	(0.4)						(100.0)			
12～	114	10		1							125	1	3	129
(%)	(91.2)	(8.0)		(0.8)							(100.0)			
合計	3,656	423	68	21	6	3	1		1		4,179	22	198	4,399
(%)	(87.5)	(10.1)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			
記載なし	102	17	2	1							122	38	2,629	2,789
(%)	(83.6)	(13.9)	(1.6)	(0.8)							(100.0)			
総計	3,758	440	70	22	6	3	1		1		4,301	60	2,827	7,188
(%)	(87.4)	(10.2)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			
平均	6.53	6.61	6.67	6.72	6.35	6.93	3.00		5.20		6.54	7.34	6.20	6.53
標準偏差	2.74	2.64	1.77	2.41	3.21	1.62					2.71	2.35	2.43	2.70

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査