

表 4482 腹膜炎罹患回数(回/年) PD透析液交換方法別(腹膜透析単独患者)

PD透析液交換方法	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
完全手動式	1,216	124	16	7	2	1			1		1,367	1	54	1,422
(%)	(89.0)	(9.1)	(1.2)	(0.5)	(0.1)	(0.1)			(0.1)		(100.0)			
バッグ交換デバイス(紫外線)	1,921	229	40	9	3	1					2,203	6	113	2,322
(%)	(87.2)	(10.4)	(1.8)	(0.4)	(0.1)	(0.0)					(100.0)			
バッグ交換デバイス(熱式無菌接合装置)	498	72	14	5	1	1					591		16	607
(%)	(84.3)	(12.2)	(2.4)	(0.8)	(0.2)	(0.2)					(100.0)			
バッグ交換デバイス(左記以外、半手動も含む)	54	5		1			1				61	1	4	66
(%)	(88.5)	(8.2)		(1.6)			(1.6)				(100.0)			
合計	3,689	430	70	22	6	3	1		1		4,222	8	187	4,417
(%)	(87.4)	(10.2)	(1.7)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			
不明	21	3									24	52		76
(%)	(87.5)	(12.5)									(100.0)			
記載なし	48	7									55		2,640	2,695
(%)	(87.3)	(12.7)									(100.0)			
総計	3,758	440	70	22	6	3	1		1		4,301	60	2,827	7,188
(%)	(87.4)	(10.2)	(1.6)	(0.5)	(0.1)	(0.1)	(0.0)		(0.0)		(100.0)			

数値下の括弧内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査