

表 4491 腹膜炎罹患回数(回/年) 透析歴(年)別 (PD・HD(F)等併用患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	123	7		1							131	6	99	236
(%)	(93.9)	(5.3)		(0.8)							(100.0)			
2年～	288	40	3	1		1					333	11	274	618
(%)	(86.5)	(12.0)	(0.9)	(0.3)		(0.3)					(100.0)			
5年～	342	29	5	3	1					1	381	2	273	656
(%)	(89.8)	(7.6)	(1.3)	(0.8)	(0.3)					(0.3)	(100.0)			
10年～	95	11	2								108		60	168
(%)	(88.0)	(10.2)	(1.9)								(100.0)			
15年～	30		2								32		19	51
(%)	(93.8)		(6.3)								(100.0)			
20年～	6										6		3	9
(%)	(100.0)										(100.0)			
25年～	3										3		1	4
(%)	(100.0)										(100.0)			
30年～	2										2		2	4
(%)	(100.0)										(100.0)			
35年～	1										1			1
(%)	(100.0)										(100.0)			
40年～														
(%)														
合計	890	87	12	5	1	1				1	997	19	731	1,747
(%)	(89.3)	(8.7)	(1.2)	(0.5)	(0.1)	(0.1)				(0.1)	(100.0)			
不明													1	1
(%)														
記載なし														
(%)														
総計	890	87	12	5	1	1				1	997	19	732	1,748
(%)	(89.3)	(8.7)	(1.2)	(0.5)	(0.1)	(0.1)				(0.1)	(100.0)			
平均	5.68	5.05	8.00	4.20	6.00	2.00				5.00	5.64	2.47	5.27	5.45
標準偏差	4.50	3.10	4.94	2.28							4.39	1.71	4.05	4.25

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査