

表 4492 腹膜炎罹患回数(回/年) 原疾患別 (PD・HD(F)等併用患者)

原疾患	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	396	40	5	3	1	1				1	447	2	278	727
(%)	(88.6)	(8.9)	(1.1)	(0.7)	(0.2)	(0.2)				(0.2)	(100.0)			
慢性腎盂腎炎	10	1									11		3	14
(%)	(90.9)	(9.1)									(100.0)			
急速進行性糸球体腎炎	6										6		4	10
(%)	(100.0)										(100.0)			
妊娠腎/妊娠中毒症	5	1	1								7		3	10
(%)	(71.4)	(14.3)	(14.3)								(100.0)			
その他の分類不能の腎炎	11										11	1	7	19
(%)	(100.0)										(100.0)			
多発性嚢胞腎	19	4									23		14	37
(%)	(82.6)	(17.4)									(100.0)			
腎硬化症	67	7	2								76	3	64	143
(%)	(88.2)	(9.2)	(2.6)								(100.0)			
悪性高血圧	14										14		8	22
(%)	(100.0)										(100.0)			
糖尿病性腎症	249	25	3	1							278	12	240	530
(%)	(89.6)	(9.0)	(1.1)	(0.4)							(100.0)			
SLE腎炎	5										5		3	8
(%)	(100.0)										(100.0)			
アミロイド腎	1										1		1	2
(%)	(100.0)										(100.0)			
痛風腎	6										6		1	7
(%)	(100.0)										(100.0)			
先天性代謝異常による腎不全	1										1			1
(%)	(100.0)										(100.0)			
腎・尿路結核														
(%)														
腎・尿路結石	1										1		1	2
(%)	(100.0)										(100.0)			
腎・尿路腫瘍	1	1									2		1	3
(%)	(50.0)	(50.0)									(100.0)			
閉塞性尿路障害	1										1		2	3
(%)	(100.0)										(100.0)			
骨髄腫													1	1
(%)														
腎形成不全	2										2		2	4
(%)	(100.0)										(100.0)			
不明	64	7	1	1							73	1	76	150
(%)	(87.7)	(9.6)	(1.4)	(1.4)							(100.0)			
再導入	9	1									10		5	15
(%)	(90.0)	(10.0)									(100.0)			
その他	22										22		18	40
(%)	(100.0)										(100.0)			
合計	890	87	12	5	1	1				1	997	19	732	1,748
(%)	(89.3)	(8.7)	(1.2)	(0.5)	(0.1)	(0.1)				(0.1)	(100.0)			
記載なし														
(%)														
総計	890	87	12	5	1	1				1	997	19	732	1,748
(%)	(89.3)	(8.7)	(1.2)	(0.5)	(0.1)	(0.1)				(0.1)	(100.0)			

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査