

表 4496 腹膜炎罹患回数(回/年) 一日透析液使用量(L/日)別 (PD・HD(F)等併用患者)

一日透析液使用量	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2未満	13										13			13
(%)	(100.0)										(100.0)			
2～	19	2	1								22			22
(%)	(86.4)	(9.1)	(4.5)								(100.0)			
4～	74	10	1								85		1	86
(%)	(87.1)	(11.8)	(1.2)								(100.0)			
6～	254	30	4	2	1	1					292		10	302
(%)	(87.0)	(10.3)	(1.4)	(0.7)	(0.3)	(0.3)					(100.0)			
8～	319	24	3	2						1	349	2	14	365
(%)	(91.4)	(6.9)	(0.9)	(0.6)						(0.3)	(100.0)			
10～	120	5	3								128	1	6	135
(%)	(93.8)	(3.9)	(2.3)								(100.0)			
12～	59	6		1							66		5	71
(%)	(89.4)	(9.1)		(1.5)							(100.0)			
合計	858	77	12	5	1	1				1	955	3	36	994
(%)	(89.8)	(8.1)	(1.3)	(0.5)	(0.1)	(0.1)				(0.1)	(100.0)			
記載なし	32	10									42	16	696	754
(%)	(76.2)	(23.8)									(100.0)			
総計	890	87	12	5	1	1				1	997	19	732	1,748
(%)	(89.3)	(8.7)	(1.2)	(0.5)	(0.1)	(0.1)				(0.1)	(100.0)			
平均	7.88	7.54	7.33	9.90	6.00	7.90				8.00	7.85	9.40	8.87	7.90
標準偏差	2.45	2.42	2.43	5.84							2.47	1.51	2.42	2.47

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査