

表 4510 出口部感染罹患回数(回/年) 透析歴(年)別 (腹膜透析患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,726	290	74	23	11	4	6			2	2,136	23	1,441	3,600
(%)	(80.8)	(13.6)	(3.5)	(1.1)	(0.5)	(0.2)	(0.3)			(0.1)	(100.0)			
2年～	1,485	254	69	21	10	8	5		1	12	1,865	49	1,287	3,201
(%)	(79.6)	(13.6)	(3.7)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.6)	(100.0)			
5年～	817	112	31	12	7	7	3		2	2	993	8	656	1,657
(%)	(82.3)	(11.3)	(3.1)	(1.2)	(0.7)	(0.7)	(0.3)		(0.2)	(0.2)	(100.0)			
10年～	159	31	7	1	1	2	1				202		138	340
(%)	(78.7)	(15.3)	(3.5)	(0.5)	(0.5)	(1.0)	(0.5)				(100.0)			
15年～	47	7	2	1							57	1	39	97
(%)	(82.5)	(12.3)	(3.5)	(1.8)							(100.0)			
20年～	12										12		8	20
(%)	(100.0)										(100.0)			
25年～	6	1									7		1	8
(%)	(85.7)	(14.3)									(100.0)			
30年～	5	1									6		4	10
(%)	(83.3)	(16.7)									(100.0)			
35年～	3										3			3
(%)	(100.0)										(100.0)			
40年～														
(%)														
合計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,574	8,936
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
不明													5	5
(%)														
記載なし														
(%)														
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	3.13	2.99	2.97	2.91	3.10	3.95	2.87		5.33	3.06	3.10	2.70	3.10	3.10
標準偏差	3.71	3.42	3.05	2.84	2.98	2.77	2.72		1.53	1.44	3.63	2.41	3.50	3.56

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査