

表 4522 出口部感染罹患回数(回/年) 身長(cm)別 (腹膜透析患者)

身長	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
130未満	39	5	3	1			1				49		13	62
(%)	(79.6)	(10.2)	(6.1)	(2.0)			(2.0)				(100.0)			
130～	9										9		2	11
(%)	(100.0)										(100.0)			
135～	38	6	1								45		17	62
(%)	(84.4)	(13.3)	(2.2)								(100.0)			
140～	140	26	5	1		2					174	2	51	227
(%)	(80.5)	(14.9)	(2.9)	(0.6)		(1.1)					(100.0)			
145～	283	38	11	7		3					342	3	125	470
(%)	(82.7)	(11.1)	(3.2)	(2.0)		(0.9)					(100.0)			
150～	528	84	18	8	3	1	3		1	1	647	10	210	867
(%)	(81.6)	(13.0)	(2.8)	(1.2)	(0.5)	(0.2)	(0.5)		(0.2)	(0.2)	(100.0)			
155～	630	109	32	7	5	1	3		1	4	792	9	248	1,049
(%)	(79.5)	(13.8)	(4.0)	(0.9)	(0.6)	(0.1)	(0.4)		(0.1)	(0.5)	(100.0)			
160～	784	120	31	8	5	5	1		1	3	958	13	359	1,330
(%)	(81.8)	(12.5)	(3.2)	(0.8)	(0.5)	(0.5)	(0.1)		(0.1)	(0.3)	(100.0)			
165～	707	113	36	12	11	4	4			1	888	17	307	1,212
(%)	(79.6)	(12.7)	(4.1)	(1.4)	(1.2)	(0.5)	(0.5)			(0.1)	(100.0)			
170～	532	97	19	8	4	4				2	666	17	221	904
(%)	(79.9)	(14.6)	(2.9)	(1.2)	(0.6)	(0.6)				(0.3)	(100.0)			
175～	203	39	13	3	1	1	1			3	264	4	98	366
(%)	(76.9)	(14.8)	(4.9)	(1.1)	(0.4)	(0.4)	(0.4)			(1.1)	(100.0)			
180～	58	13	2								73		14	87
(%)	(79.5)	(17.8)	(2.7)								(100.0)			
185～	6		1								7	1	5	13
(%)	(85.7)		(14.3)								(100.0)			
190～	4										4			4
(%)	(100.0)										(100.0)			
合計	3,961	650	172	55	29	21	13		3	14	4,918	76	1,670	6,664
(%)	(80.5)	(13.2)	(3.5)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	299	46	11	3			2			2	363	5	1,909	2,277
(%)	(82.4)	(12.7)	(3.0)	(0.8)			(0.6)			(0.6)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	160.35	161.00	160.65	160.45	163.55	160.81	156.69		157.67	164.86	160.47	163.38	160.67	160.56
標準偏差	11.75	11.18	13.15	10.66	6.71	10.35	17.22		6.11	8.80	11.70	9.03	11.37	11.59

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査