

表 4523 出口部感染罹患回数(回/年) 体重(kg)別 (腹膜透析患者)

体重	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
20未満	28	3	2								33		5	38
(%)	(84.8)	(9.1)	(6.1)								(100.0)			
20～	16	2	2	1							21		2	23
(%)	(76.2)	(9.5)	(9.5)	(4.8)							(100.0)			
30～	108	25	6	2							141	1	30	172
(%)	(76.6)	(17.7)	(4.3)	(1.4)							(100.0)			
40～	608	88	27	8	2	3	4			1	741	1	162	904
(%)	(82.1)	(11.9)	(3.6)	(1.1)	(0.3)	(0.4)	(0.5)			(0.1)	(100.0)			
50～	1,074	148	37	12	4	8	5		1	1	1,290	11	290	1,591
(%)	(83.3)	(11.5)	(2.9)	(0.9)	(0.3)	(0.6)	(0.4)		(0.1)	(0.1)	(100.0)			
60～	1,059	186	47	17	7	6	2		2	5	1,331	14	271	1,616
(%)	(79.6)	(14.0)	(3.5)	(1.3)	(0.5)	(0.5)	(0.2)		(0.2)	(0.4)	(100.0)			
70～	537	80	20	8	5	2				4	656	9	146	811
(%)	(81.9)	(12.2)	(3.0)	(1.2)	(0.8)	(0.3)				(0.6)	(100.0)			
80～	188	30	10	2	4	1	2			2	239	1	58	298
(%)	(78.7)	(12.6)	(4.2)	(0.8)	(1.7)	(0.4)	(0.8)			(0.8)	(100.0)			
90～	62	11	5		1					1	80	1	23	104
(%)	(77.5)	(13.8)	(6.3)		(1.3)					(1.3)	(100.0)			
100～	13	3		1							17	1	2	20
(%)	(76.5)	(17.6)		(5.9)							(100.0)			
110～	6	1	1								8		2	10
(%)	(75.0)	(12.5)	(12.5)								(100.0)			
120～	6		1								7		2	9
(%)	(85.7)		(14.3)								(100.0)			
合計	3,705	577	158	51	23	20	13		3	14	4,564	39	993	5,596
(%)	(81.2)	(12.6)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	555	119	25	7	6	1	2			2	717	42	2,586	3,345
(%)	(77.4)	(16.6)	(3.5)	(1.0)	(0.8)	(0.1)	(0.3)			(0.3)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	60.53	60.76	61.06	61.25	69.02	60.03	58.92		61.27	69.54	60.65	65.51	61.01	60.74
標準偏差	13.94	13.58	15.88	14.76	12.53	9.91	13.29		4.60	12.86	13.96	12.68	13.63	13.90

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査