

表 4524 出口部感染罹患回数(回/年) Body Mass Index別（腹膜透析患者）

Body Mass Index	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
14未満	22		1								23		4	27
(%)	(95.7)		(4.3)								(100.0)			
14～	45	13	1								59	1	11	71
(%)	(76.3)	(22.0)	(1.7)								(100.0)			
16～	150	26	8	5		1	1				191		37	228
(%)	(78.5)	(13.6)	(4.2)	(2.6)		(0.5)	(0.5)				(100.0)			
18～	427	68	23	3	1	3	2			1	528	2	97	627
(%)	(80.9)	(12.9)	(4.4)	(0.6)	(0.2)	(0.6)	(0.4)			(0.2)	(100.0)			
20～	744	121	27	8	5	4	3			2	914	5	221	1,140
(%)	(81.4)	(13.2)	(3.0)	(0.9)	(0.5)	(0.4)	(0.3)			(0.2)	(100.0)			
22～	818	126	36	14	2	4	2		1	4	1,007	11	243	1,261
(%)	(81.2)	(12.5)	(3.6)	(1.4)	(0.2)	(0.4)	(0.2)		(0.1)	(0.4)	(100.0)			
24～	628	95	22	11	4	4	1		2		767	9	132	908
(%)	(81.9)	(12.4)	(2.9)	(1.4)	(0.5)	(0.5)	(0.1)		(0.3)		(100.0)			
26～	373	59	18	5	6	4	2			2	469	2	90	561
(%)	(79.5)	(12.6)	(3.8)	(1.1)	(1.3)	(0.9)	(0.4)			(0.4)	(100.0)			
28～	199	31	9	2	2					2	245	3	54	302
(%)	(81.2)	(12.7)	(3.7)	(0.8)	(0.8)					(0.8)	(100.0)			
30～	140	25	7	1	3		1			3	180	3	37	220
(%)	(77.8)	(13.9)	(3.9)	(0.6)	(1.7)		(0.6)			(1.7)	(100.0)			
合計	3,546	564	152	49	23	20	12		3	14	4,383	36	926	5,345
(%)	(80.9)	(12.9)	(3.5)	(1.1)	(0.5)	(0.5)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	714	132	31	9	6	1	3			2	898	45	2,653	3,596
(%)	(79.5)	(14.7)	(3.5)	(1.0)	(0.7)	(0.1)	(0.3)			(0.2)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	23.13	23.14	23.15	23.23	25.32	22.92	23.02		24.63	25.64	23.15	24.14	23.06	23.15
標準偏差	3.59	3.73	3.73	3.42	3.71	2.75	4.04		0.85	4.77	3.62	3.66	3.49	3.60

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査