

表 4525 出口部感染罹患回数(回/年) BUN(mg/dL)別(腹膜透析患者)

BUN	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
20未満	12	2									14	1	14	29
(%)	(85.7)	(14.3)									(100.0)			
20～	111	18	3	1	1	1			1		136	1	46	183
(%)	(81.6)	(13.2)	(2.2)	(0.7)	(0.7)	(0.7)			(0.7)		(100.0)			
30～	436	55	20	3	5	2				1	522	6	144	672
(%)	(83.5)	(10.5)	(3.8)	(0.6)	(1.0)	(0.4)				(0.2)	(100.0)			
40～	917	149	43	12	7	3	4			5	1,140	6	246	1,392
(%)	(80.4)	(13.1)	(3.8)	(1.1)	(0.6)	(0.3)	(0.4)			(0.4)	(100.0)			
50～	1,069	152	51	20	3	3	1		1	4	1,304	13	296	1,613
(%)	(82.0)	(11.7)	(3.9)	(1.5)	(0.2)	(0.2)	(0.1)		(0.1)	(0.3)	(100.0)			
60～	742	131	19	9	6	7	6		1	2	923	6	221	1,150
(%)	(80.4)	(14.2)	(2.1)	(1.0)	(0.7)	(0.8)	(0.7)		(0.1)	(0.2)	(100.0)			
70～	359	53	20	6	3	1	1			1	444	8	97	549
(%)	(80.9)	(11.9)	(4.5)	(1.4)	(0.7)	(0.2)	(0.2)			(0.2)	(100.0)			
80～	157	25	3	2	2	2	2				193	1	51	245
(%)	(81.3)	(13.0)	(1.6)	(1.0)	(1.0)	(1.0)	(1.0)				(100.0)			
90～	48	9	2	1		1				1	62	2	25	89
(%)	(77.4)	(14.5)	(3.2)	(1.6)		(1.6)				(1.6)	(100.0)			
100～	21	4	1			1					27	1	12	40
(%)	(77.8)	(14.8)	(3.7)			(3.7)					(100.0)			
110～	2	1	1								4		3	7
(%)	(50.0)	(25.0)	(25.0)								(100.0)			
120～	6										6		1	7
(%)	(100.0)										(100.0)			
合計	3,880	599	163	54	27	21	14		3	14	4,775	45	1,156	5,976
(%)	(81.3)	(12.5)	(3.4)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	380	97	20	4	2		1			2	506	36	2,423	2,965
(%)	(75.1)	(19.2)	(4.0)	(0.8)	(0.4)		(0.2)			(0.4)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	54.79	55.33	54.31	56.04	53.37	60.24	62.57		47.00	56.00	54.89	56.89	54.87	54.90
標準偏差	15.34	15.02	14.78	13.60	16.26	19.16	12.27		21.00	13.68	15.28	19.59	17.77	15.82

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査