

表 4528 出口部感染罹患回数(回/年) CRP濃度(mg/dL)別(腹膜透析患者)

CRP濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
0.2未満	2,030	302	89	28	13	6	7		3	6	2,484	16	567	3,067
(%)	(81.7)	(12.2)	(3.6)	(1.1)	(0.5)	(0.2)	(0.3)		(0.1)	(0.2)	(100.0)			
0.2～	660	96	26	9	5	3	2			1	802	18	175	995
(%)	(82.3)	(12.0)	(3.2)	(1.1)	(0.6)	(0.4)	(0.2)			(0.1)	(100.0)			
0.5～	298	37	8	3	2	2	1			2	353	7	89	449
(%)	(84.4)	(10.5)	(2.3)	(0.8)	(0.6)	(0.6)	(0.3)			(0.6)	(100.0)			
1.0～	178	34	13	2	1	4				3	235	2	71	308
(%)	(75.7)	(14.5)	(5.5)	(0.9)	(0.4)	(1.7)				(1.3)	(100.0)			
2.0～	113	28	8	2		2	2				155	2	45	202
(%)	(72.9)	(18.1)	(5.2)	(1.3)		(1.3)	(1.3)				(100.0)			
4.0～	49	9	1		1						60	1	22	83
(%)	(81.7)	(15.0)	(1.7)		(1.7)						(100.0)			
6.0～	20	7	1	1							29		11	40
(%)	(69.0)	(24.1)	(3.4)	(3.4)							(100.0)			
8.0～	13	1			1		1				16		8	24
(%)	(81.3)	(6.3)			(6.3)		(6.3)				(100.0)			
10.0～	24	3	2								29		5	34
(%)	(82.8)	(10.3)	(6.9)								(100.0)			
15.0～	4	2									6		1	7
(%)	(66.7)	(33.3)									(100.0)			
合計	3,389	519	148	45	23	17	13		3	12	4,169	46	994	5,209
(%)	(81.3)	(12.4)	(3.6)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	871	177	35	13	6	4	2			4	1,112	35	2,585	3,732
(%)	(78.3)	(15.9)	(3.1)	(1.2)	(0.5)	(0.4)	(0.2)			(0.4)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	0.58	0.78	0.66	0.46	0.82	0.81	1.21		0.05	0.48	0.61	0.51	0.72	0.63
標準偏差	1.58	2.57	1.62	1.07	2.09	0.92	2.45		0.06	0.50	1.73	0.91	1.65	1.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査