

表 4530 出口部感染罹患回数(回/年) リン濃度(mg/dL)別 (腹膜透析患者)

リン濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
3未満	65	8	8	2	1	2					86	1	39	126
(%)	(75.6)	(9.3)	(9.3)	(2.3)	(1.2)	(2.3)					(100.0)			
3.0～	173	29	6	2	4				1		215	1	54	270
(%)	(80.5)	(13.5)	(2.8)	(0.9)	(1.9)				(0.5)		(100.0)			
3.5～	330	52	19	2	3	2	1			1	410	6	109	525
(%)	(80.5)	(12.7)	(4.6)	(0.5)	(0.7)	(0.5)	(0.2)			(0.2)	(100.0)			
4.0～	516	62	17	9	3	3	4			3	617	4	149	770
(%)	(83.6)	(10.0)	(2.8)	(1.5)	(0.5)	(0.5)	(0.6)			(0.5)	(100.0)			
4.5～	613	108	23	9	2	3	1			2	761	4	156	921
(%)	(80.6)	(14.2)	(3.0)	(1.2)	(0.3)	(0.4)	(0.1)			(0.3)	(100.0)			
5.0～	630	101	29	5	5	2	2				774	3	171	948
(%)	(81.4)	(13.0)	(3.7)	(0.6)	(0.6)	(0.3)	(0.3)				(100.0)			
5.5～	557	90	20	2	3	4	2			3	681	6	152	839
(%)	(81.8)	(13.2)	(2.9)	(0.3)	(0.4)	(0.6)	(0.3)			(0.4)	(100.0)			
6.0～	91	14	3	2							110	1	27	138
(%)	(82.7)	(12.7)	(2.7)	(1.8)							(100.0)			
6.1～	272	41	15	9		3	3			3	346	7	124	477
(%)	(78.6)	(11.8)	(4.3)	(2.6)		(0.9)	(0.9)			(0.9)	(100.0)			
6.5～	260	41	10	5	1				2		319	1	79	399
(%)	(81.5)	(12.9)	(3.1)	(1.6)	(0.3)				(0.6)		(100.0)			
7.0～	145	21	7	2	2	1				1	179	2	31	212
(%)	(81.0)	(11.7)	(3.9)	(1.1)	(1.1)	(0.6)				(0.6)	(100.0)			
7.5～	99	10	3	1	1	1	1				116	6	27	149
(%)	(85.3)	(8.6)	(2.6)	(0.9)	(0.9)	(0.9)	(0.9)				(100.0)			
8.0～	42	7	3	3							55	2	14	71
(%)	(76.4)	(12.7)	(5.5)	(5.5)							(100.0)			
8.5～	19	7		1							27		11	38
(%)	(70.4)	(25.9)		(3.7)							(100.0)			
9.0～	64	3	1		2					1	71	1	17	89
(%)	(90.1)	(4.2)	(1.4)		(2.8)					(1.4)	(100.0)			
合計	3,876	594	164	54	27	21	14		3	14	4,767	45	1,160	5,972
(%)	(81.3)	(12.5)	(3.4)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	384	102	19	4	2		1			2	514	36	2,419	2,969
(%)	(74.7)	(19.8)	(3.7)	(0.8)	(0.4)		(0.2)			(0.4)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	5.26	5.22	5.12	5.44	5.14	5.01	5.19		5.63	5.55	5.25	5.65	5.22	5.25
標準偏差	1.34	1.22	1.35	1.44	1.81	1.49	1.20		2.19	1.47	1.33	1.64	1.41	1.35

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査