

表 4533 出口部感染罹患回数(回/年) ヘモグロビン濃度(g/dL)別 (腹膜透析患者)

ヘモグロビン 濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
5未満	5	2									7			7
(%)	(71.4)	(28.6)									(100.0)			
5～	9	2	1								12	1	1	14
(%)	(75.0)	(16.7)	(8.3)								(100.0)			
6～	17	3	1								21	1	6	28
(%)	(81.0)	(14.3)	(4.8)								(100.0)			
7～	69	11	1	1						1	83	4	37	124
(%)	(83.1)	(13.3)	(1.2)	(1.2)						(1.2)	(100.0)			
8～	208	21	13	2	2	1	1				248	6	90	344
(%)	(83.9)	(8.5)	(5.2)	(0.8)	(0.8)	(0.4)	(0.4)				(100.0)			
9～	597	73	20	6	3	6	1			1	707	11	185	903
(%)	(84.4)	(10.3)	(2.8)	(0.8)	(0.4)	(0.8)	(0.1)			(0.1)	(100.0)			
10～	1,082	198	44	16	8	8	4		1	6	1,367	11	328	1,706
(%)	(79.2)	(14.5)	(3.2)	(1.2)	(0.6)	(0.6)	(0.3)		(0.1)	(0.4)	(100.0)			
11～	1,159	176	50	13	5	3	7			3	1,416	7	286	1,709
(%)	(81.9)	(12.4)	(3.5)	(0.9)	(0.4)	(0.2)	(0.5)			(0.2)	(100.0)			
12～	491	86	24	13	5	3	1		2	3	628	4	164	796
(%)	(78.2)	(13.7)	(3.8)	(2.1)	(0.8)	(0.5)	(0.2)		(0.3)	(0.5)	(100.0)			
13～	162	17	7	2	3						191	1	45	237
(%)	(84.8)	(8.9)	(3.7)	(1.0)	(1.6)						(100.0)			
14～	51	7	1		1						60		19	79
(%)	(85.0)	(11.7)	(1.7)		(1.7)						(100.0)			
合計	3,850	596	162	53	27	21	14		3	14	4,740	46	1,161	5,947
(%)	(81.2)	(12.6)	(3.4)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	410	100	21	5	2		1			2	541	35	2,418	2,994
(%)	(75.8)	(18.5)	(3.9)	(0.9)	(0.4)		(0.2)			(0.4)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	10.83	10.87	10.84	11.07	11.37	10.48	10.80		11.50	10.94	10.84	10.00	10.72	10.81
標準偏差	1.40	1.41	1.40	1.28	1.57	1.03	1.05		0.95	1.41	1.40	1.61	1.47	1.42

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査