

表 4534 出口部感染罹患回数(回/年) 総コレステロール濃度(mg/dL)別（腹膜透析患者）

総コレステロール濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
100未満	29	3	3	2		1	1				39		11	50
(%)	(74.4)	(7.7)	(7.7)	(5.1)		(2.6)	(2.6)				(100.0)			
100～	106	14	8	1	1						130	2	28	160
(%)	(81.5)	(10.8)	(6.2)	(0.8)	(0.8)						(100.0)			
120～	290	51	7	5	2	4	1		1	1	362		86	448
(%)	(80.1)	(14.1)	(1.9)	(1.4)	(0.6)	(1.1)	(0.3)		(0.3)	(0.3)	(100.0)			
140～	541	89	25	5	2	4	1			2	669	6	131	806
(%)	(80.9)	(13.3)	(3.7)	(0.7)	(0.3)	(0.6)	(0.1)			(0.3)	(100.0)			
160～	616	86	21	7	1	2	3			1	737	10	118	865
(%)	(83.6)	(11.7)	(2.8)	(0.9)	(0.1)	(0.3)	(0.4)			(0.1)	(100.0)			
180～	599	90	28	6	2	1	1		2		729	6	98	833
(%)	(82.2)	(12.3)	(3.8)	(0.8)	(0.3)	(0.1)	(0.1)		(0.3)		(100.0)			
200～	393	55	20	5	1	2					476	4	84	564
(%)	(82.6)	(11.6)	(4.2)	(1.1)	(0.2)	(0.4)					(100.0)			
220～	237	31	5	1	1	1	1			1	278	2	38	318
(%)	(85.3)	(11.2)	(1.8)	(0.4)	(0.4)	(0.4)	(0.4)			(0.4)	(100.0)			
240～	115	14	5		1						135		24	159
(%)	(85.2)	(10.4)	(3.7)		(0.7)						(100.0)			
260～	105	17	3	2		1					128	2	19	149
(%)	(82.0)	(13.3)	(2.3)	(1.6)		(0.8)					(100.0)			
合計	3,031	450	125	34	11	16	8		3	5	3,683	32	637	4,352
(%)	(82.3)	(12.2)	(3.4)	(0.9)	(0.3)	(0.4)	(0.2)		(0.1)	(0.1)	(100.0)			
記載なし	1,229	246	58	24	18	5	7			11	1,598	49	2,942	4,589
(%)	(76.9)	(15.4)	(3.6)	(1.5)	(1.1)	(0.3)	(0.4)			(0.7)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	180.37	178.46	177.89	173.35	172.36	172.00	158.00		162.33	164.60	179.84	180.41	174.14	179.01
標準偏差	40.35	39.36	42.92	44.41	41.77	65.11	39.79		35.85	40.20	40.48	38.20	40.67	40.54

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査