

表 4536 出口部感染罹患回数(回/年) Non-HDL-C濃度(mg/dL)別(腹膜透析患者)

Non-HDL-C濃度	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
50未満	13		1	1							15		3	18
(%)	(86.7)		(6.7)	(6.7)							(100.0)			
50～	70	11	5	1	1	2	1				91		23	114
(%)	(76.9)	(12.1)	(5.5)	(1.1)	(1.1)	(2.2)	(1.1)				(100.0)			
70～	234	41	9	4	1	2			1	1	293		58	351
(%)	(79.9)	(14.0)	(3.1)	(1.4)	(0.3)	(0.7)			(0.3)	(0.3)	(100.0)			
90～	475	62	17	6	3	2	2			2	569	2	89	660
(%)	(83.5)	(10.9)	(3.0)	(1.1)	(0.5)	(0.4)	(0.4)			(0.4)	(100.0)			
110～	550	88	23	6	1	3	1				672	8	117	797
(%)	(81.8)	(13.1)	(3.4)	(0.9)	(0.1)	(0.4)	(0.1)				(100.0)			
130～	475	65	19	4	1	1	2		2		569	1	92	662
(%)	(83.5)	(11.4)	(3.3)	(0.7)	(0.2)	(0.2)	(0.4)		(0.4)		(100.0)			
150～	305	48	17	4	2	2					378	3	55	436
(%)	(80.7)	(12.7)	(4.5)	(1.1)	(0.5)	(0.5)					(100.0)			
170～	191	26	7		2					1	227	1	35	263
(%)	(84.1)	(11.5)	(3.1)		(0.9)					(0.4)	(100.0)			
190～	85	12	3	3							103	2	15	120
(%)	(82.5)	(11.7)	(2.9)	(2.9)							(100.0)			
210～	65	10	2			1					78	1	11	90
(%)	(83.3)	(12.8)	(2.6)			(1.3)					(100.0)			
合計	2,463	363	103	29	11	13	6		3	4	2,995	18	498	3,511
(%)	(82.2)	(12.1)	(3.4)	(1.0)	(0.4)	(0.4)	(0.2)		(0.1)	(0.1)	(100.0)			
記載なし	1,797	333	80	29	18	8	9			12	2,286	63	3,081	5,430
(%)	(78.6)	(14.6)	(3.5)	(1.3)	(0.8)	(0.3)	(0.4)			(0.5)	(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	129.01	128.98	128.56	122.62	125.82	126.54	112.67		128.00	120.00	128.86	143.78	125.57	128.47
標準偏差	37.78	37.40	39.09	41.02	39.22	71.24	29.33		34.64	47.79	37.97	37.84	38.11	38.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査