

表 4541 出口部感染罹患回数(回/年) 使用透析液の種類別(腹膜透析患者)

使用透析液の種類	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
1.5%液のみ	1,717	281	73	33	10	4	7			6	2,131	10	134	2,275
(%)	(80.6)	(13.2)	(3.4)	(1.5)	(0.5)	(0.2)	(0.3)			(0.3)	(100.0)			
1.5%液と2.5%液の併用	583	96	25	8	7	2	2		1	2	726	9	52	787
(%)	(80.3)	(13.2)	(3.4)	(1.1)	(1.0)	(0.3)	(0.3)		(0.1)	(0.3)	(100.0)			
2.5%液のみ	134	20	7		1						162	2	13	177
(%)	(82.7)	(12.3)	(4.3)		(0.6)						(100.0)			
4.25%液のみ(イコデキストリンなし)	8										8			8
(%)	(100.0)										(100.0)			
イコデキストリン単独(ブドウ糖使用なし)	48	4	2			1					55		2	57
(%)	(87.3)	(7.3)	(3.6)			(1.8)					(100.0)			
1.5%+イコデキストリン	920	146	45	15	6	6	2		1	4	1,145	4	73	1,222
(%)	(80.3)	(12.8)	(3.9)	(1.3)	(0.5)	(0.5)	(0.2)		(0.1)	(0.3)	(100.0)			
1.5%+2.5%+イコデキストリン	415	80	17	1	5	2	3		1	3	527	1	31	559
(%)	(78.7)	(15.2)	(3.2)	(0.2)	(0.9)	(0.4)	(0.6)		(0.2)	(0.6)	(100.0)			
2.5%+イコデキストリン	336	57	13	1		6	1			1	415	3	23	441
(%)	(81.0)	(13.7)	(3.1)	(0.2)		(1.4)	(0.2)			(0.2)	(100.0)			
4.25%液+イコデキストリン	4	2									6			6
(%)	(66.7)	(33.3)									(100.0)			
合計	4,165	686	182	58	29	21	15		3	16	5,175	29	328	5,532
(%)	(80.5)	(13.3)	(3.5)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
不明	9	2									11	52	2	65
(%)	(81.8)	(18.2)									(100.0)			
記載なし	86	8	1								95		3,249	3,344
(%)	(90.5)	(8.4)	(1.1)								(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査