

表 4542 出口部感染罹患回数(回/年) 一日透析液使用量(L/日)別(腹膜透析患者)

一日透析液使用量	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2未満	88	13	4	1	2	1					109		10	119
(%)	(80.7)	(11.9)	(3.7)	(0.9)	(1.8)	(0.9)					(100.0)			
2～	306	46	13	7	1	1	1			1	376	2	15	393
(%)	(81.4)	(12.2)	(3.5)	(1.9)	(0.3)	(0.3)	(0.3)			(0.3)	(100.0)			
4～	790	172	35	13	8	1	4				1,023		49	1,072
(%)	(77.2)	(16.8)	(3.4)	(1.3)	(0.8)	(0.1)	(0.4)				(100.0)			
6～	1,368	232	63	19	8	7	6			6	1,709	4	82	1,795
(%)	(80.0)	(13.6)	(3.7)	(1.1)	(0.5)	(0.4)	(0.4)			(0.4)	(100.0)			
8～	1,091	145	41	9	8	10	2		2	6	1,314	18	74	1,406
(%)	(83.0)	(11.0)	(3.1)	(0.7)	(0.6)	(0.8)	(0.2)		(0.2)	(0.5)	(100.0)			
10～	327	43	12	8	2	1	1			2	396	2	12	410
(%)	(82.6)	(10.9)	(3.0)	(2.0)	(0.5)	(0.3)	(0.3)			(0.5)	(100.0)			
12～	146	29	13	1			1		1	1	192	1	8	201
(%)	(76.0)	(15.1)	(6.8)	(0.5)			(0.5)		(0.5)	(0.5)	(100.0)			
合計	4,116	680	181	58	29	21	15		3	16	5,119	27	250	5,396
(%)	(80.4)	(13.3)	(3.5)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
記載なし	144	16	2								162	54	3,329	3,545
(%)	(88.9)	(9.9)	(1.2)								(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
平均	6.82	6.59	6.90	6.42	6.23	7.01	6.71		9.33	8.10	6.79	7.66	6.64	6.78
標準偏差	2.77	2.47	2.74	2.50	2.34	2.10	2.19		2.31	2.12	2.72	2.11	2.57	2.71

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査