

表 4547 出口部感染罹患回数(回/年) PD透析液交換方法別(腹膜透析患者)

PD透析液交換方法	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
完全手動式	1,375	225	56	11	6	5	3			7	1,688	2	70	1,760
(%)	(81.5)	(13.3)	(3.3)	(0.7)	(0.4)	(0.3)	(0.2)			(0.4)	(100.0)			
バッグ交換デバイス(紫外線)	2,186	368	98	34	17	12	10		2	5	2,732	7	136	2,875
(%)	(80.0)	(13.5)	(3.6)	(1.2)	(0.6)	(0.4)	(0.4)		(0.1)	(0.2)	(100.0)			
バッグ交換デバイス(熱式無菌接合装置)	561	83	23	11	5	1	1		1	3	689	28	717	
(%)	(81.4)	(12.0)	(3.3)	(1.6)	(0.7)	(0.1)	(0.1)		(0.1)	(0.4)	(100.0)			
バッグ交換デバイス(左記以外、半手動も含む)	54	9	5	2	1	1	1			1	74	1	5	80
(%)	(73.0)	(12.2)	(6.8)	(2.7)	(1.4)	(1.4)	(1.4)			(1.4)	(100.0)			
合計	4,176	685	182	58	29	19	15		3	16	5,183	10	239	5,432
(%)	(80.6)	(13.2)	(3.5)	(1.1)	(0.6)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			
不明	30	3									33	71	3	107
(%)	(90.9)	(9.1)									(100.0)			
記載なし	54	8	1			2					65		3,337	3,402
(%)	(83.1)	(12.3)	(1.5)			(3.1)					(100.0)			
総計	4,260	696	183	58	29	21	15		3	16	5,281	81	3,579	8,941
(%)	(80.7)	(13.2)	(3.5)	(1.1)	(0.5)	(0.4)	(0.3)		(0.1)	(0.3)	(100.0)			

数値下の括弧内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査