

表 4558 出口部感染罹患回数(回/年) 使用透析液の種類別(腹膜透析単独患者)

使用透析液の種類	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
1.5%液のみ	1,575	268	66	27	9	4	7			6	1,962	8	121	2,091
(%)	(80.3)	(13.7)	(3.4)	(1.4)	(0.5)	(0.2)	(0.4)			(0.3)	(100.0)			
1.5%液と2.5%液の併用	419	67	17	6	5		1		1	2	518	7	40	565
(%)	(80.9)	(12.9)	(3.3)	(1.2)	(1.0)		(0.2)		(0.2)	(0.4)	(100.0)			
2.5%液のみ	97	13	5		1						116	2	8	126
(%)	(83.6)	(11.2)	(4.3)		(0.9)						(100.0)			
4.25%液のみ(イコデキストリンなし)	7										7			7
(%)	(100.0)										(100.0)			
イコデキストリン単独(ブドウ糖使用なし)	42	4	2		1						49		2	51
(%)	(85.7)	(8.2)	(4.1)		(2.0)						(100.0)			
1.5%+イコデキストリン	759	114	38	10	5	3	2		4	935	3	67	1,005	
(%)	(81.2)	(12.2)	(4.1)	(1.1)	(0.5)	(0.3)	(0.2)		(0.4)	(100.0)				
1.5%+2.5%+イコデキストリン	268	52	12	1	4	1	2		2	342	1	23	366	
(%)	(78.4)	(15.2)	(3.5)	(0.3)	(1.2)	(0.3)	(0.6)		(0.6)	(100.0)				
2.5%+イコデキストリン	223	40	9	1		4				277	3	14	294	
(%)	(80.5)	(14.4)	(3.2)	(0.4)		(1.4)				(100.0)				
4.25%液+イコデキストリン	1	2								3				3
(%)	(33.3)	(66.7)								(100.0)				
合計	3,391	560	149	45	24	13	12		1	14	4,209	24	275	4,508
(%)	(80.6)	(13.3)	(3.5)	(1.1)	(0.6)	(0.3)	(0.3)		(0.0)	(0.3)	(100.0)			
不明	9										9	37	2	48
(%)	(100.0)										(100.0)			
記載なし	65	5	1								71		2,561	2,632
(%)	(91.5)	(7.0)	(1.4)								(100.0)			
総計	3,465	565	150	45	24	13	12		1	14	4,289	61	2,838	7,188
(%)	(80.8)	(13.2)	(3.5)	(1.0)	(0.6)	(0.3)	(0.3)		(0.0)	(0.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査