

表 4559 出口部感染罹患回数(回/年) 一日透析液使用量(L/日)別 (腹膜透析単独患者)

一日透析液使用量	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2未満	80	10	4	1		1					96		10	106
(%)	(83.3)	(10.4)	(4.2)	(1.0)		(1.0)					(100.0)			
2～	286	46	13	6	1	1	1				354	2	15	371
(%)	(80.8)	(13.0)	(3.7)	(1.7)	(0.3)	(0.3)	(0.3)				(100.0)			
4～	726	158	33	12	8		2				939		47	986
(%)	(77.3)	(16.8)	(3.5)	(1.3)	(0.9)		(0.2)				(100.0)			
6～	1,138	187	55	15	8	5	6			5	1,419	3	71	1,493
(%)	(80.2)	(13.2)	(3.9)	(1.1)	(0.6)	(0.4)	(0.4)			(0.4)	(100.0)			
8～	805	108	30	6	6	5	1		1	6	968	16	57	1,041
(%)	(83.2)	(11.2)	(3.1)	(0.6)	(0.6)	(0.5)	(0.1)		(0.1)	(0.6)	(100.0)			
10～	224	29	3	5	1	1	1			2	266	1	6	273
(%)	(84.2)	(10.9)	(1.1)	(1.9)	(0.4)	(0.4)	(0.4)			(0.8)	(100.0)			
12～	95	17	11				1			1	125	1	3	129
(%)	(76.0)	(13.6)	(8.8)				(0.8)			(0.8)	(100.0)			
合計	3,354	555	149	45	24	13	12		1	14	4,167	23	209	4,399
(%)	(80.5)	(13.3)	(3.6)	(1.1)	(0.6)	(0.3)	(0.3)		(0.0)	(0.3)	(100.0)			
記載なし	111	10	1								122	38	2,629	2,789
(%)	(91.0)	(8.2)	(0.8)								(100.0)			
総計	3,465	565	150	45	24	13	12		1	14	4,289	61	2,838	7,188
(%)	(80.8)	(13.2)	(3.5)	(1.0)	(0.6)	(0.3)	(0.3)		(0.0)	(0.3)	(100.0)			
平均	6.57	6.32	6.54	5.96	6.30	6.78	6.88		8.00	8.51	6.54	7.50	6.24	6.53
標準偏差	2.78	2.37	2.75	2.29	1.78	2.56	2.32			1.72	2.72	2.12	2.42	2.70

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査