

表 4572 出口部感染罹患回数(回/年) 透析歴(年)別 (PD・HD(F)等併用患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	107	17	3	1		1					129	6	101	236
(%)	(82.9)	(13.2)	(2.3)	(0.8)		(0.8)					(100.0)			
2年～	251	51	17	7	1	2	1			2	332	11	275	618
(%)	(75.6)	(15.4)	(5.1)	(2.1)	(0.3)	(0.6)	(0.3)			(0.6)	(100.0)			
5年～	315	38	9	5	4	3	1		2		377	3	276	656
(%)	(83.6)	(10.1)	(2.4)	(1.3)	(1.1)	(0.8)	(0.3)		(0.5)		(100.0)			
10年～	82	18	4			2	1				107		61	168
(%)	(76.6)	(16.8)	(3.7)			(1.9)	(0.9)				(100.0)			
15年～	26	6									32		19	51
(%)	(81.3)	(18.8)									(100.0)			
20年～	6										6		3	9
(%)	(100.0)										(100.0)			
25年～	3										3		1	4
(%)	(100.0)										(100.0)			
30年～	2										2		2	4
(%)	(100.0)										(100.0)			
35年～	1										1			1
(%)	(100.0)										(100.0)			
40年～														
(%)														
合計	793	130	33	13	5	8	3		2	2	989	20	738	1,747
(%)	(80.2)	(13.1)	(3.3)	(1.3)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			
不明													1	1
(%)														
記載なし														
(%)														
総計	793	130	33	13	5	8	3		2	2	989	20	739	1,748
(%)	(80.2)	(13.1)	(3.3)	(1.3)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			
平均	5.73	5.49	5.03	3.62	5.60	5.25	6.33		6.00	4.00	5.65	2.65	5.27	5.45
標準偏差	4.55	4.02	3.40	1.71	1.14	3.54	3.21		1.41		4.40	1.84	4.05	4.25

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査