

表 4573 出口部感染罹患回数(回/年) 原疾患別 (PD・HD(F)等併用患者)

原疾患	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	356	59	15	5	2	4	1		1		443	3	281	727
(%)	(80.4)	(13.3)	(3.4)	(1.1)	(0.5)	(0.9)	(0.2)		(0.2)		(100.0)			
慢性腎盂腎炎	11										11		3	14
(%)	(100.0)										(100.0)			
急速進行性糸球体腎炎	2	3		1							6		4	10
(%)	(33.3)	(50.0)		(16.7)							(100.0)			
妊娠腎/妊娠中毒症	6	1									7		3	10
(%)	(85.7)	(14.3)									(100.0)			
その他の分類不能の腎炎	8	1		1		1					11	1	7	19
(%)	(72.7)	(9.1)		(9.1)		(9.1)					(100.0)			
多発性嚢胞腎	20		2			1					23		14	37
(%)	(87.0)		(8.7)			(4.3)					(100.0)			
腎硬化症	62	10	2			1			1		76	3	64	143
(%)	(81.6)	(13.2)	(2.6)			(1.3)			(1.3)		(100.0)			
悪性高血圧	12	2									14		8	22
(%)	(85.7)	(14.3)									(100.0)			
糖尿病性腎症	213	45	10	3	1	1	2			1	276	12	242	530
(%)	(77.2)	(16.3)	(3.6)	(1.1)	(0.4)	(0.4)	(0.7)			(0.4)	(100.0)			
SLE腎炎	4										4		4	8
(%)	(100.0)										(100.0)			
アミロイド腎	1										1		1	2
(%)	(100.0)										(100.0)			
痛風腎	5	1									6		1	7
(%)	(83.3)	(16.7)									(100.0)			
先天性代謝異常による腎不全	1										1			1
(%)	(100.0)										(100.0)			
腎・尿路結核														
(%)														
腎・尿路結石	1										1		1	2
(%)	(100.0)										(100.0)			
腎・尿路腫瘍	2										2		1	3
(%)	(100.0)										(100.0)			
閉塞性尿路障害	1										1		2	3
(%)	(100.0)										(100.0)			
骨髄腫													1	1
(%)														
腎形成不全	1	1									2		2	4
(%)	(50.0)	(50.0)									(100.0)			
不明	60	5	4	3							72	1	77	150
(%)	(83.3)	(6.9)	(5.6)	(4.2)							(100.0)			
再導入	9	1									10		5	15
(%)	(90.0)	(10.0)									(100.0)			
その他	18	1			2					1	22		18	40
(%)	(81.8)	(4.5)			(9.1)					(4.5)	(100.0)			
合計	793	130	33	13	5	8	3		2	2	989	20	739	1,748
(%)	(80.2)	(13.1)	(3.3)	(1.3)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			
記載なし														
(%)														
総計	793	130	33	13	5	8	3		2	2	989	20	739	1,748
(%)	(80.2)	(13.1)	(3.3)	(1.3)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			

数値下のかつこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査