

表 4576 出口部感染罹患回数(回/年) 使用透析液の種類別 (PD・HD(F)等併用患者)

使用透析液の種類	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
1.5%液のみ	142	13	7	6	1						169	2	13	184
(%)	(84.0)	(7.7)	(4.1)	(3.6)	(0.6)						(100.0)			
1.5%液と2.5%液の併用	164	28	8	2	2	2	1				207	2	12	221
(%)	(79.2)	(13.5)	(3.9)	(1.0)	(1.0)	(1.0)	(0.5)				(100.0)			
2.5%液のみ	37	7	2								46		5	51
(%)	(80.4)	(15.2)	(4.3)								(100.0)			
4.25%液のみ(イコデキストリンなし)	1										1			1
(%)	(100.0)										(100.0)			
イコデキストリン単独(ブドウ糖使用なし)	6										6			6
(%)	(100.0)										(100.0)			
1.5%+イコデキストリン	161	32	7	5	1	3			1		210	1	6	217
(%)	(76.7)	(15.2)	(3.3)	(2.4)	(0.5)	(1.4)			(0.5)		(100.0)			
1.5%+2.5%+イコデキストリン	147	28	5		1	1	1		1	1	185		8	193
(%)	(79.5)	(15.1)	(2.7)		(0.5)	(0.5)	(0.5)		(0.5)	(0.5)	(100.0)			
2.5%+イコデキストリン	111	17	4			2	1			1	136		9	145
(%)	(81.6)	(12.5)	(2.9)			(1.5)	(0.7)			(0.7)	(100.0)			
4.25%液+イコデキストリン	3										3			3
(%)	(100.0)										(100.0)			
合計	772	125	33	13	5	8	3		2	2	963	5	53	1,021
(%)	(80.2)	(13.0)	(3.4)	(1.3)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			
不明		2									2		15	
(%)		(100.0)									(100.0)			17
記載なし	21	3									24		686	710
(%)	(87.5)	(12.5)									(100.0)			
総計	793	130	33	13	5	8	3		2	2	989	20	739	1,748
(%)	(80.2)	(13.1)	(3.3)	(1.3)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査