

表 4577 出口部感染罹患回数(回/年) 一日透析液使用量(L/日)別 (PD・HD(F)等併用患者)

一日透析液使用量	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2未満	8	3			2						13			13
(%)	(61.5)	(23.1)			(15.4)						(100.0)			
2～	20			1						1	22			22
(%)	(90.9)			(4.5)						(4.5)	(100.0)			
4～	64	14	2	1		1	2				84		2	86
(%)	(76.2)	(16.7)	(2.4)	(1.2)		(1.2)	(2.4)				(100.0)			
6～	230	45	8	4		2				1	290	1	11	302
(%)	(79.3)	(15.5)	(2.8)	(1.4)		(0.7)				(0.3)	(100.0)			
8～	286	37	11	3	2	5	1		1		346	2	17	365
(%)	(82.7)	(10.7)	(3.2)	(0.9)	(0.6)	(1.4)	(0.3)		(0.3)		(100.0)			
10～	102	13	9	3	1						128	1	6	135
(%)	(79.7)	(10.2)	(7.0)	(2.3)	(0.8)						(100.0)			
12～	50	12	2	1					1		66		5	71
(%)	(75.8)	(18.2)	(3.0)	(1.5)					(1.5)		(100.0)			
合計	760	124	32	13	5	8	3		2	2	949	4	41	994
(%)	(80.1)	(13.1)	(3.4)	(1.4)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			
記載なし	33	6	1								40	16	698	754
(%)	(82.5)	(15.0)	(2.5)								(100.0)			
総計	793	130	33	13	5	8	3		2	2	989	20	739	1,748
(%)	(80.2)	(13.1)	(3.3)	(1.3)	(0.5)	(0.8)	(0.3)		(0.2)	(0.2)	(100.0)			
平均	7.87	7.77	8.59	7.99	5.94	7.38	6.00		10.00	5.25	7.86	8.55	8.68	7.90
標準偏差	2.47	2.53	1.97	2.66	4.48	1.03	1.73		2.83	3.18	2.48	2.10	2.36	2.47

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査