

表 124 2015年導入患者, 性別, 原疾患別

原疾患	男性	女性	合計	記載なし	総計
慢性糸球体腎炎	4,014	2,218	6,232		6,232
(%)	(16.1)	(18.8)	(16.9)		(16.9)
慢性腎盂腎炎	127	129	256		256
(%)	(0.5)	(1.1)	(0.7)		(0.7)
急速進行性糸球体腎炎	271	192	463		463
(%)	(1.1)	(1.6)	(1.3)		(1.3)
妊娠腎/妊娠中毒症	1	29	30		30
(%)	(0.0)	(0.2)	(0.1)		(0.1)
その他の分類不能の腎炎	86	55	141		141
(%)	(0.3)	(0.5)	(0.4)		(0.4)
多発性嚢胞腎	542	397	939		939
(%)	(2.2)	(3.4)	(2.6)		(2.6)
腎硬化症	3,540	1,685	5,225		5,225
(%)	(14.2)	(14.3)	(14.2)		(14.2)
悪性高血圧	198	98	296		296
(%)	(0.8)	(0.8)	(0.8)		(0.8)
糖尿病性腎症	11,680	4,392	16,072		16,072
(%)	(46.7)	(37.3)	(43.7)		(43.7)
SLE腎炎	106	163	269		269
(%)	(0.4)	(1.4)	(0.7)		(0.7)
アミロイド腎	56	56	112		112
(%)	(0.2)	(0.5)	(0.3)		(0.3)
痛風腎	64	19	83		83
(%)	(0.3)	(0.2)	(0.2)		(0.2)
先天性代謝異常による腎不全	13	6	19		19
(%)	(0.1)	(0.1)	(0.1)		(0.1)
腎・尿路結核	7	2	9		9
(%)	(0.0)	(0.0)	(0.0)		(0.0)
腎・尿路結石	51	25	76		76
(%)	(0.2)	(0.2)	(0.2)		(0.2)
腎・尿路腫瘍	148	29	177		177
(%)	(0.6)	(0.2)	(0.5)		(0.5)
閉塞性尿路障害	74	30	104		104
(%)	(0.3)	(0.3)	(0.3)		(0.3)
骨髄腫	92	50	142		142
(%)	(0.4)	(0.4)	(0.4)		(0.4)
腎形成不全	35	16	51		51
(%)	(0.1)	(0.1)	(0.1)		(0.1)
不明	2,880	1,593	4,473		4,473
(%)	(11.5)	(13.5)	(12.2)		(12.2)
再導入	129	87	216		216
(%)	(0.5)	(0.7)	(0.6)		(0.6)
その他	892	519	1,411		1,411
(%)	(3.6)	(4.4)	(3.8)		(3.8)
合計	25,006	11,790	36,796		36,796
(%)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし	1		1		1
総計	25,007	11,790	36,797		36,797

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査