

表 128 2015年導入患者，原疾患別，平均年齢

原疾患	人数	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	6,232		6,232	68.77	14.37
(%)	(16.9)		(16.9)		
慢性腎盂腎炎	256		256	67.56	15.69
(%)	(0.7)		(0.7)		
急速進行性糸球体腎炎	463		463	73.12	12.13
(%)	(1.3)		(1.3)		
妊娠腎／妊娠中毒症	30		30	62.60	13.63
(%)	(0.1)		(0.1)		
その他分類不能の腎炎	141		141	67.42	16.08
(%)	(0.4)		(0.4)		
多発性嚢胞腎	939		939	62.50	13.69
(%)	(2.6)		(2.6)		
腎硬化症	5,225		5,225	75.33	11.31
(%)	(14.2)		(14.2)		
悪性高血圧	296		296	63.61	17.74
(%)	(0.8)		(0.8)		
糖尿病性腎症	16,070	2	16,072	67.29	12.21
(%)	(43.7)	(40.0)	(43.7)		
SLE腎炎	269		269	63.76	14.35
(%)	(0.7)		(0.7)		
アミロイド腎	112		112	69.72	10.61
(%)	(0.3)		(0.3)		
痛風腎	83		83	62.70	14.38
(%)	(0.2)		(0.2)		
先天性代謝異常による腎不全	19		19	48.37	23.20
(%)	(0.1)		(0.1)		
腎・尿路結核	9		9	81.00	8.49
(%)	(0.0)		(0.0)		
腎・尿路結石	76		76	71.75	12.14
(%)	(0.2)		(0.2)		
腎・尿路腫瘍	177		177	72.74	10.38
(%)	(0.5)		(0.5)		
閉塞性尿路障害	104		104	68.98	13.56
(%)	(0.3)		(0.3)		
骨髄腫	142		142	71.73	9.87
(%)	(0.4)		(0.4)		
腎形成不全	51		51	45.47	26.79
(%)	(0.1)		(0.1)		
不明	4,470	3	4,473	72.04	13.38
(%)	(12.1)	(60.0)	(12.2)		
移植後再導入	216		216	55.67	15.07
(%)	(0.6)		(0.6)		
その他	1,411		1,411	69.68	14.96
(%)	(3.8)		(3.8)		
合計	36,791	5	36,796	69.20	13.39
(%)	(100.0)	(100.0)	(100.0)		
記載なし	1		1	74.00	
総計	36,792	5	36,797	69.20	13.39

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査