

表 138 2015年導入患者死亡原因分類(透析期間との関係)

死因	3ヶ月未満	3ヶ月～	6ヶ月～	1 年 ～	2 年 ～	5 年 ～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
心不全	267	162	105								534			534
(%)	(24.3)	(25.0)	(25.5)								(24.8)			(24.7)
脳血管障害	47	50	27								124			124
(%)	(4.3)	(7.7)	(6.6)								(5.8)			(5.7)
感染症	315	149	93								557			557
(%)	(28.7)	(23.0)	(22.6)								(25.8)			(25.8)
消化管出血	22	8	7								37			37
(%)	(2.0)	(1.2)	(1.7)								(1.7)			(1.7)
悪性腫瘍	98	91	46								235		1	236
(%)	(8.9)	(14.1)	(11.2)								(10.9)		(33.3)	(10.9)
悪液質/尿毒症	57	27	22								106			106
(%)	(5.2)	(4.2)	(5.4)								(4.9)			(4.9)
心筋梗塞	35	8	18								61			61
(%)	(3.2)	(1.2)	(4.4)								(2.8)			(2.8)
カリウム中毒/頓死	15	11	5								31			31
(%)	(1.4)	(1.7)	(1.2)								(1.4)			(1.4)
肝硬変症	16	9	8								33			33
(%)	(1.5)	(1.4)	(1.9)								(1.5)			(1.5)
自殺/拒否	8	4	3								15			15
(%)	(0.7)	(0.6)	(0.7)								(0.7)			(0.7)
腸閉塞	14	7	3								24			24
(%)	(1.3)	(1.1)	(0.7)								(1.1)			(1.1)
肺血栓/肺塞栓	1	2									3			3
(%)	(0.1)	(0.3)									(0.1)			(0.1)
災害・事故死	5		3								8			8
(%)	(0.5)		(0.7)								(0.4)			(0.4)
その他	126	69	36								231			231
(%)	(11.5)	(10.7)	(8.8)								(10.7)			(10.7)
不明	72	50	35								157		2	159
(%)	(6.6)	(7.7)	(8.5)								(7.3)		(66.7)	(7.4)
合計	1,098	647	411								2,156		3	2,159
(%)	(100.0)	(100.0)	(100.0)								(100.0)		(100.0)	(100.0)
記載なし		1	2								3			3
総計	1,098	648	413								2,159		3	2,162

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査