

表 144 2015年導入患者死亡原因分類(年齢別、原疾患別)原疾患=その他分類不能の腎炎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差									
心不全 (%)															1 (25.0)	1 (100.0)	2 (50.0)	1 (100.0)	1 (100.0)	1 (100.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	2 (25.0)	6.36
脳血管障害 (%)																																			
感染症 (%)																																			
消化管出血 (%)																																			
悪性腫瘍 (%)																																			
悪液質/尿毒症 (%)																																			
心筋梗塞 (%)																																			
カリウム中毒/頓死 (%)																																			
肝硬変症 (%)																																			
自殺/拒否 (%)																																			
腸閉塞 (%)																																			
肺血栓/肺塞栓 (%)																																			
災害・事故死 (%)																																			
その他 (%)																																			
不明 (%)																																			
合計 (%)										1 (100.0)				1 (100.0)		4 (100.0)	1 (100.0)	1 (100.0)			8 (100.0)			8 (100.0)		8 (100.0)		74.88	11.42						
記載なし																																			
総計										1				1		4	1	1			8			8		74.88	11.42								

数値下の括弧内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査