

表 145 2015年導入患者死亡原因分類(年齢別、原疾患別)原疾患=多発性囊胞腎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)										1 (100.0)				2 (28.6)		2 (50.0)	1 (100.0)		1 (50.0)		7 (26.9)			7 (26.9)	72.86	12.19	
脳血管障害 (%)											1 (33.3)										1 (3.8)			1 (3.8)	62.00		
感染症 (%)												1 (14.3)									2 (7.7)			2 (7.7)	77.00	11.31	
消化管出血 (%)												1 (33.3)									2 (7.7)			2 (7.7)	73.00	18.38	
悪性腫瘍 (%)													2 (28.6)	1 (25.0)	1 (25.0)							4 (15.4)			4 (15.4)	70.75	5.25
悪液質/尿毒症 (%)																					2 (7.7)			2 (7.7)	70.00	2.83	
心筋梗塞 (%)														1 (14.3)	1 (25.0)							2 (7.7)			2 (7.7)	72.00	21.21
カリウム中毒/頓死 (%)																											
肝硬変症 (%)																											
自殺/拒否 (%)																											
腸閉塞 (%)																											
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)																					3 (11.5)			3 (11.5)	65.67	4.04	
不明 (%)																					2 (7.7)			2 (7.7)	82.00	12.73	
合計 (%)										1 (100.0)	1 (100.0)	3 (100.0)	7 (100.0)	4 (100.0)	4 (100.0)	1 (100.0)	3 (100.0)	2 (100.0)	26 (100.0)				26 (100.0)	72.19	10.21		
記載なし																											
総計										1	1	3	7	4	4	1	3	2	26				26	72.19	10.21		

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査