

表 147 2015年導入患者死亡原因分類(年齢別、原疾患別)原疾患=悪性高血圧

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)						1 (100.0)											1 (50.0)	2 (100.0)	4 (44.4)			4 (44.4)		74.25	31.67	
脳血管障害 (%)																	1 (100.0)	1 (50.0)	1 (100.0)	1 (11.1)	1 (11.1)	1 (11.1)	1 (11.1)	1 (11.1)	79.00	
感染症 (%)																	1 (100.0)	1 (100.0)	1 (100.0)	1 (11.1)	1 (11.1)	1 (11.1)	1 (11.1)	1 (11.1)	66.00	
消化管出血 (%)																										
悪性腫瘍 (%)																										
悪液質/尿毒症 (%)																										
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																										
不明 (%)																										
合計 (%)						1 (100.0)								1 (100.0)		2 (100.0)	1 (100.0)	2 (100.0)	1 (100.0)	2 (100.0)	1 (100.0)	9 (100.0)		9 (100.0)	77.00	20.43
記載なし																										
総計						1								1		2	1	2	2		9			9	77.00	20.43

数値下の括弧内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査