

表 150 2015年導入患者死亡原因分類(年齢別、原疾患別)原疾患=アミロイド腎

| 死因 | 5歳未満 | 5歳～ | 10歳～ | 15歳～ | 20歳～ | 25歳～ | 30歳～ | 35歳～ | 40歳～ | 45歳～ | 50歳～ | 55歳～ | 60歳～ | 65歳～ | 70歳～ | 75歳～ | 80歳～ | 85歳～ | 90歳～ | 95歳～ | 合計 | 不明 | 記載なし | 総計 | 平均 | 標準偏差 | | |
|------------------|------|-----|------|------|------|------|------|------|------|------|------|------|------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|-------------|-------------|---------------|-------------|-------------|---------------|-------|------|
| 心不全 (%) | | | | | | | | | | | | | | 3 (75.0) | 1 (16.7) | 1 (100.0) | | | | | 5 (25.0) | | | 5 (25.0) | 75.00 | 4.95 | | |
| 脳血管障害 (%) | | | | | | | | | | | | | | | 2 (33.3) | | | | | | 2 (10.0) | | | 2 (10.0) | 77.00 | 2.83 | | |
| 感染症 (%) | | | | | | | | | | | | | | 2 (66.7) | 1 (33.3) | 1 (25.0) | | | | | 5 (25.0) | | | 5 (25.0) | 70.00 | 11.40 | | |
| 消化管出血 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 悪性腫瘍 (%) | | | | | | | | | | | | | | 1 (100.0) | | | 1 (16.7) | | | | | 3 (15.0) | | | 3 (15.0) | 73.67 | 14.98 | |
| 悪液質/尿毒症 (%) | | | | | | | | | | | | | | | | 1 (16.7) | | | | | 1 (5.0) | | | 1 (5.0) | 77.00 | | | |
| 心筋梗塞 (%) | | | | | | | | | | | | | | | | | | | | | 1 (5.0) | | | 1 (5.0) | 76.00 | | | |
| カリウム中毒/頓死 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 肝硬変症 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 自殺/拒否 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 腸閉塞 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 肺血栓/肺塞栓 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 災害・事故死 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| その他 (%) | | | | | | | | | | | | | | 1 (33.3) | 2 (66.7) | | | | | | | 3 (15.0) | | | 3 (15.0) | 66.00 | 2.00 | |
| 不明 (%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 合計 (%) | | | | | | | | | | | | | | 1 (100.0) | 3 (100.0) | 3 (100.0) | 4 (100.0) | 6 (100.0) | 1 (100.0) | 2 (100.0) | | | 20 (100.0) | | | 20 (100.0) | 72.55 | 8.43 |
| 記載なし | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 総計 | | | | | | | | | | | | | | 1 | 3 | 3 | 4 | 6 | 1 | 2 | | | 20 | | 20 | 72.55 | 8.43 | |

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査