

表 154 2015年導入患者死亡原因分類(年齢別, 原疾患別)原疾患＝腎・尿路結石

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)																	1 (100.0)				1 (33.3)			1 (33.3)	84.00	
脳血管障害 (%)																										
感染症 (%)																		1 (100.0)			1 (33.3)			1 (33.3)	88.00	
消化管出血 (%)																										
悪性腫瘍 (%)																										
悪液質/尿毒症 (%)																										
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																1 (100.0)					1 (33.3)			1 (33.3)	75.00	
不明 (%)																										
合計 (%)																1 (100.0)	1 (100.0)	1 (100.0)			3 (100.0)			3 (100.0)	82.33	6.66
記載なし																										
総計																1	1	1			3			3	82.33	6.66

数値下のかっこ内は列方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査