

表 155 2015年導入患者死亡原因分類(年齢別、原疾患別)原疾患=腎・尿路腫瘍

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差		
心不全 (%)															1 (20.0)	1 (10.0)	1 (33.3)				3		3 (13.0)		83.00	5.57		
脳血管障害 (%)														1 (50.0)							1 (4.3)			1 (4.3)		70.00		
感染症 (%)															3 (30.0)						3 (13.0)		3 (13.0)		80.33	0.58		
消化管出血 (%)																												
悪性腫瘍 (%)														1 (50.0)	1 (50.0)	4 (80.0)	6 (60.0)	2 (66.7)				14 (60.9)			14 (60.9)		78.57	6.05
悪液質/尿毒症 (%)																												
心筋梗塞 (%)																												
カリウム中毒/頓死 (%)																												
肝硬変症 (%)																												
自殺/拒否 (%)																												
腸閉塞 (%)																												
肺血栓/肺塞栓 (%)																												
災害・事故死 (%)																												
その他 (%)														1 (50.0)							1 (4.3)			1 (4.3)		64.00		
不明 (%)															1 (100.0)						1 (4.3)			1 (4.3)		67.00		
合計 (%)											2 (100.0)	1 (100.0)	2 (100.0)	5 (100.0)	10 (100.0)	3 (100.0)				23 (100.0)			23 (100.0)		77.87	6.79		
記載なし																												
総計											2	1	2	5	10	3				23			23		77.87	6.79		

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査