

表 157 2015年導入患者死亡原因分類(年齢別、原疾患別)原疾患=骨髄腫

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)																					4			4	79.25	4.03
脳血管障害 (%)																									(10.5)	
感染症 (%)																					5			5	66.60	14.74
消化管出血 (%)																									(13.2)	
悪性腫瘍 (%)																					27			27	72.93	10.80
悪液質/尿毒症 (%)																									(71.1)	
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																					1			1	87.00	
不明 (%)																					(2.6)			(2.6)		
合計 (%)										1 (100.0)	2 (100.0)	1 (100.0)	4 (100.0)	5 (100.0)	6 (100.0)	7 (100.0)	6 (100.0)	5 (100.0)	1 (100.0)	38 (100.0)				38	73.18	11.04
記載なし																										
総計										1	2	1	4	5	6	7	6	5	1	38				38	73.18	11.04

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査