

表 160 2015年導入患者死亡原因分類(年齢別, 原疾患別)原疾患＝移植後再導入

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)										1 (100.0)				1 (100.0)	1 (33.3)						3 (30.0)			3 (30.0)	62.67	13.32
脳血管障害 (%)																	1 (33.3)				1 (10.0)			1 (10.0)	81.00	
感染症 (%)											1 (100.0)	1 (100.0)			1 (33.3)		1 (33.3)				4 (40.0)			4 (40.0)	65.75	13.05
消化管出血 (%)																										
悪性腫瘍 (%)																										
悪液質/尿毒症 (%)																										
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)															1 (33.3)						1 (10.0)			1 (10.0)	70.00	
肝硬変症 (%)																										
自殺/拒否 (%)																	1 (33.3)				1 (10.0)			1 (10.0)	84.00	
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																										
不明 (%)																										
合計 (%)										1 (100.0)	1 (100.0)	1 (100.0)		1 (100.0)	3 (100.0)		3 (100.0)				10 (100.0)			10 (100.0)	68.60	12.46
記載なし																										
総計										1	1	1		1	3		3				10			10	68.60	12.46

数値下のかっこ内は列方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査