

表 218 2015年末患者、年齢、原疾患別人数(女性)

原疾患	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2	1	1	10	24	62	176	412	942	1,446	2,169	2,864	4,765	6,691	5,823	4,975	3,780	2,265	686	103	37,197			37,197	68.10	12.32
(%)	(8.7)	(7.7)	(7.1)	(23.3)	(27.6)	(31.0)	(31.9)	(35.3)	(36.9)	(37.2)	(38.4)	(36.7)	(37.3)	(35.1)	(34.1)	(31.2)	(27.8)	(27.2)	(25.5)	(25.8)	(33.3)			(33.2)		
慢性腎盂腎炎					2	7	15	35	50	86	110	118	170	266	240	192	172	79	27	4	1,573			1,573	66.68	13.47
(%)					(2.3)	(3.5)	(2.7)	(3.0)	(2.0)	(2.2)	(1.9)	(1.5)	(1.3)	(1.4)	(1.4)	(1.2)	(1.3)	(0.9)	(1.0)	(1.0)	(1.4)			(1.4)		
急速進行性糸球体腎炎			2	1	5	5	7	13	24	43	39	69	118	165	188	180	197	105	29	1	1,191			1,191	70.30	13.28
(%)			(14.3)	(2.3)	(5.7)	(2.5)	(1.3)	(1.1)	(0.9)	(1.1)	(0.7)	(0.9)	(0.9)	(0.9)	(1.1)	(1.1)	(1.5)	(1.3)	(1.1)	(0.3)	(1.1)			(1.1)		
妊娠腎/妊娠中毒症							7	7	30	81	119	162	250	364	242	161	84	23	3		1,533			1,533	65.10	10.19
(%)							(1.3)	(0.6)	(1.2)	(2.1)	(2.1)	(2.1)	(2.0)	(1.9)	(1.4)	(1.0)	(0.6)	(0.3)	(0.1)		(1.4)			(1.4)		
その他の分類不能の腎炎	1		1	1	5	10	12	20	27	49	41	49	70	64	67	64	64	29	8	2	584			584	63.30	16.19
(%)	(4.3)		(7.1)	(2.3)	(5.7)	(5.0)	(2.2)	(1.7)	(1.1)	(1.3)	(0.7)	(0.6)	(0.5)	(0.3)	(0.4)	(0.4)	(0.5)	(0.3)	(0.3)	(0.5)	(0.5)			(0.5)		
多発性囊胞腎	2			1	3	4	16	33	89	176	368	493	833	1,050	802	628	377	157	34	7	5,073			5,073	66.55	10.85
(%)	(8.7)			(2.3)	(3.4)	(2.0)	(2.9)	(2.8)	(3.5)	(4.5)	(6.5)	(6.3)	(6.5)	(5.5)	(4.7)	(3.9)	(2.8)	(1.9)	(1.3)	(1.8)	(4.5)			(4.5)		
腎硬化症				2	5	6	16	32	84	139	185	300	555	1,090	1,283	1,710	2,120	1,734	661	138	10,060			10,060	76.48	11.19
(%)				(4.7)	(5.7)	(3.0)	(2.9)	(2.7)	(3.3)	(3.8)	(3.3)	(3.8)	(4.4)	(5.7)	(7.5)	(10.7)	(15.6)	(20.8)	(24.5)	(34.5)	(9.0)			(9.0)		
悪性高血圧					1	2	7	15	31	50	54	62	89	122	99	116	103	59	31	6	847			847	68.07	14.20
(%)					(1.1)	(1.0)	(1.3)	(1.3)	(1.2)	(1.3)	(1.0)	(0.8)	(0.7)	(0.6)	(0.6)	(0.7)	(0.8)	(0.7)	(1.2)	(1.5)	(0.8)			(0.8)		
糖尿病性腎症	1		1	2	3	20	103	292	695	1,066	1,663	2,475	4,051	6,585	5,969	5,413	4,258	2,230	596	47	35,470	1		35,471	69.46	11.51
(%)	(4.3)		(7.1)	(4.7)	(3.4)	(10.0)	(18.7)	(25.0)	(27.2)	(27.4)	(29.4)	(31.7)	(31.8)	(34.6)	(34.9)	(33.9)	(31.4)	(26.8)	(22.1)	(11.8)	(31.7)	(50.0)		(31.7)		
SLE腎炎				1	1	10	19	43	93	135	150	189	261	241	162	100	72	40	14	1	1,532			1,532	61.22	12.91
(%)				(2.3)	(1.1)	(5.0)	(3.4)	(3.7)	(3.6)	(3.5)	(2.7)	(2.4)	(2.0)	(1.3)	(0.9)	(0.6)	(0.5)	(0.5)	(0.5)	(0.3)	(1.4)			(1.4)		
アミロイド腎								1	3	5	10	14	31	55	48	48	33	12	2		262			262	70.23	9.73
(%)								(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.3)	(0.2)	(0.1)	(0.1)		(0.2)			(0.2)		
痛風腎								1	2	12	8	7	15	23	18	25	12	11	2		136			136	68.51	12.40
(%)								(0.1)	(0.1)	(0.3)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.1)	(0.1)	(0.1)		(0.1)			(0.1)		
先天性代謝異常による腎不全	1				5	2	10	10	14	13	9	7	11	13	8	4	4	4			115			115	52.40	17.43
(%)	(4.3)				(5.7)	(1.0)	(1.8)	(0.9)	(0.5)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)			(0.1)			(0.1)		
腎・尿路結核										1	2	2	8	15	25	9	20	6		1	89			89	73.37	8.80
(%)										(0.0)	(0.0)	(0.0)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)		(0.3)	(0.1)			(0.1)		
腎・尿路結石							1	1	4		11	10	18	29	33	27	41	20	4	3	202			202	72.63	11.76
(%)							(0.2)	(0.1)	(0.2)		(0.2)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)	(0.3)	(0.2)	(0.1)	(0.8)	(0.2)			(0.2)		
腎・尿路腫瘍								5	2	4	5	5	20	30	37	41	38	16	6		209			209	72.42	11.28
(%)								(0.4)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.2)	(0.2)		(0.2)			(0.2)		
閉塞性尿路障害	1					6	8	14	13	13	18	15	22	36	27	27	28	15	5		248			248	63.54	16.83
(%)	(4.3)					(3.0)	(1.5)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)		(0.2)			(0.2)		
骨髄腫									2	3	5	5	9	22	21	12	19	7	2		107			107	70.92	10.90
(%)									(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)		(0.1)			(0.1)		
腎形成不全	5	4	4	8	7	11	21	27	36	23	16	14	13	13	13	15	7	9	3		249			249	48.05	20.77
(%)	(21.7)	(30.8)	(28.6)	(18.6)	(8.0)	(5.5)	(3.8)	(2.3)	(1.4)	(0.8)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)		(0.2)			(0.2)		
不明	1		1	7	11	12	59	108	219	331	411	650	1,044	1,639	1,538	1,770	1,723	1,246	491	71	11,332	1		11,333	71.75	12.89
(%)	(4.3)		(7.1)	(16.3)	(12.6)	(6.0)	(10.7)	(9.3)	(8.6)	(8.5)	(7.3)	(8.3)	(8.2)	(8.6)	(9.0)	(11.1)	(12.7)	(15.0)	(18.2)	(17.8)	(10.1)	(50.0)		(10.1)		
再導入				2	10	11	17	37	70	69	97	113	113	108	38	24	19	16	3	2	749			749	56.57	13.60
(%)				(4.7)	(11.5)	(5.5)	(3.1)	(3.2)	(2.7)	(1.8)	(1.7)	(1.4)	(0.9)	(0.6)	(0.2)	(0.2)	(0.1)	(0.2)	(0.1)	(0.5)	(0.7)			(0.7)		
その他	9	8	4	8	5	32	57	61	123	146	164	191	292	433	409	426	408	234	88	14	3,112			3,112	67.30	15.92
(%)	(39.1)	(61.5)	(28.6)	(18.6)	(5.7)	(16.0)	(10.3)	(5.2)	(4.8)	(3.8)	(2.9)	(2.4)	(2.3)	(2.3)	(2.4)	(2.7)	(3.0)	(2.8)	(3.3)	(3.5)	(2.8)			(2.8)		
合計	23	13	14	43	87	200	551	1,167	2,553	3,891	5,654	7,815	12,759	19,055	17,091	15,968	13,582	8,317	2,695	400	111,878	2		111,872	69.28	12.58
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)			(100.0)		
記載なし												1	1	1	1	1	3				8			8	72.50	11.10
総計	23	13	14	43	87	200	551	1,167	2,553	3,891	5,654	7,815	12,759	19,055	17,091	15,968	13,582	8,317	2,695	400	111,878	2		111,880	69.28	12.58

数値下のかっこ内は列方向の合計に対する％です。
 (C)Japanese Society for Dialysis Therapy
 2015年調査