

表 219 2015年末患者，透析歴，原疾患別人数

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	12,359	15,982	19,909	15,345	11,873	8,008	4,839	2,885	1,630	484	93,314	32	1	93,347	11.25	9.39
(%)	(17.8)	(20.3)	(25.6)	(38.3)	(55.1)	(66.6)	(71.9)	(73.7)	(77.8)	(78.4)	(29.8)	(8.2)	(100.0)	(29.8)		
慢性腎盂腎炎	448	570	636	464	305	206	145	105	42	12	2,933	2		2,935	10.40	9.34
(%)	(0.6)	(0.7)	(0.8)	(1.2)	(1.4)	(1.7)	(2.2)	(2.7)	(2.0)	(1.9)	(0.9)	(0.5)		(0.9)		
急速進行性糸球体腎炎	689	715	565	274	112	72	25	15	6	4	2,477	1		2,478	5.79	6.45
(%)	(1.0)	(0.9)	(0.7)	(0.7)	(0.5)	(0.6)	(0.4)	(0.4)	(0.3)	(0.6)	(0.8)	(0.3)		(0.8)		
妊娠腎/妊娠中毒症	74	123	221	239	260	232	189	132	60	15	1,545			1,545	17.17	10.21
(%)	(0.1)	(0.2)	(0.3)	(0.6)	(1.2)	(1.9)	(2.8)	(3.4)	(2.9)	(2.4)	(0.5)			(0.5)		
その他の分類不能の腎炎	270	290	310	189	108	107	70	33	23	5	1,405			1,405	9.63	9.31
(%)	(0.4)	(0.4)	(0.4)	(0.5)	(0.5)	(0.9)	(1.0)	(0.8)	(1.1)	(0.8)	(0.4)			(0.4)		
多発性嚢胞腎	1,921	2,427	2,884	1,985	1,123	549	240	83	33	9	11,254	2		11,256	8.25	7.15
(%)	(2.8)	(3.1)	(3.7)	(5.0)	(5.2)	(4.6)	(3.6)	(2.1)	(1.6)	(1.5)	(3.6)	(0.5)		(3.6)		
腎硬化症	9,630	9,249	7,115	2,522	840	282	105	37	13	3	29,796	9		29,805	4.42	4.73
(%)	(13.9)	(11.8)	(9.1)	(6.3)	(3.9)	(2.3)	(1.6)	(0.9)	(0.6)	(0.5)	(9.5)	(2.3)		(9.5)		
悪性高血圧	581	649	691	332	211	93	49	5	4	3	2,618			2,618	6.87	6.60
(%)	(0.8)	(0.8)	(0.9)	(0.8)	(1.0)	(0.8)	(0.7)	(0.1)	(0.2)	(0.5)	(0.8)			(0.8)		
糖尿病性腎症	31,350	36,655	34,377	13,088	3,662	867	180	40	16	4	120,239	39		120,278	4.89	4.49
(%)	(45.2)	(46.6)	(44.2)	(32.7)	(17.0)	(7.2)	(2.7)	(1.0)	(0.8)	(0.6)	(38.4)	(9.9)		(38.4)		
SLE腎炎	460	474	467	322	226	126	83	51	16	2	2,227			2,227	8.81	8.45
(%)	(0.7)	(0.6)	(0.6)	(0.8)	(1.0)	(1.0)	(1.2)	(1.3)	(0.8)	(0.3)	(0.7)			(0.7)		
アミロイド腎	156	134	104	41	16	6	4		1		462			462	4.70	5.39
(%)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.0)	(0.1)		(0.0)		(0.1)			(0.1)		
痛風腎	156	206	262	184	133	80	34	11	5	2	1,073	2		1,075	9.47	7.85
(%)	(0.2)	(0.3)	(0.3)	(0.5)	(0.6)	(0.7)	(0.5)	(0.3)	(0.2)	(0.3)	(0.3)	(0.5)		(0.3)		
先天性代謝異常による腎不全	38	52	64	43	31	19	16	6	5		274			274	10.51	9.03
(%)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)	(0.2)		(0.1)			(0.1)		
腎・尿路結核	23	36	33	28	26	21	10	14	3	1	195			195	12.57	10.33
(%)	(0.0)	(0.0)	(0.0)	(0.1)	(0.1)	(0.2)	(0.1)	(0.4)	(0.1)	(0.2)	(0.1)			(0.1)		
腎・尿路結石	131	126	156	75	48	29	10	2	4		581			581	7.39	7.21
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)	(0.2)		(0.2)			(0.2)		
腎・尿路腫瘍	292	263	221	83	29	15	3	3	1		910			910	4.81	5.29
(%)	(0.4)	(0.3)	(0.3)	(0.2)	(0.1)	(0.1)	(0.0)	(0.1)	(0.0)		(0.3)			(0.3)		
閉塞性尿路障害	161	165	162	93	66	39	28	12	2	2	730			730	8.22	8.19
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.4)	(0.3)	(0.1)	(0.3)	(0.2)			(0.2)		
骨髄腫	153	91	33	11	1		1	1			291			291	2.47	3.60
(%)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)		(0.0)	(0.0)			(0.1)			(0.1)		
腎形成不全	99	112	122	105	85	63	35	25	11	3	660			660	11.40	9.64
(%)	(0.1)	(0.1)	(0.2)	(0.3)	(0.4)	(0.5)	(0.5)	(0.6)	(0.5)	(0.5)	(0.2)			(0.2)		
不明	7,826	7,820	7,089	3,463	1,766	813	392	271	130	37	29,607	290		29,897	6.32	6.85
(%)	(11.3)	(9.9)	(9.1)	(8.6)	(8.2)	(6.8)	(5.8)	(6.9)	(6.2)	(6.0)	(9.5)	(74.0)		(9.5)		
再導入	379	403	456	295	197	152	147	100	38	16	2,183			2,183	10.90	10.15
(%)	(0.5)	(0.5)	(0.6)	(0.7)	(0.9)	(1.3)	(2.2)	(2.6)	(1.8)	(2.6)	(0.7)			(0.7)		
その他	2,175	2,089	1,922	877	437	251	125	85	53	15	8,029	15		8,044	6.40	7.22
(%)	(3.1)	(2.7)	(2.5)	(2.2)	(2.0)	(2.1)	(1.9)	(2.2)	(2.5)	(2.4)	(2.6)	(3.8)		(2.6)		
合計	69,371	78,631	77,799	40,058	21,555	12,030	6,730	3,916	2,096	617	312,803	392	1	313,196	7.31	7.55
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		
記載なし	8	7	2	2		1		1			21			21	5.33	7.90
総計	69,379	78,638	77,801	40,060	21,555	12,031	6,730	3,917	2,096	617	312,824	392	1	313,217	7.31	7.55

数値下のかつこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査