

表 220 2015年末患者，透析歴，原疾患別人数(男性)

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	7,927	10,141	12,257	9,041	6,876	4,515	2,638	1,559	905	271	56,130	19	1	56,150	10.78	9.23
(%)	(16.9)	(19.1)	(24.2)	(36.9)	(55.3)	(68.5)	(74.1)	(77.4)	(81.1)	(79.5)	(27.9)	(7.4)	(100.0)	(27.9)		
慢性腎盂腎炎	220	298	284	215	121	88	56	45	25	8	1,360	2		1,362	10.00	9.41
(%)	(0.5)	(0.6)	(0.6)	(0.9)	(1.0)	(1.3)	(1.6)	(2.2)	(2.2)	(2.3)	(0.7)	(0.8)		(0.7)		
急速進行性糸球体腎炎	364	398	279	134	52	30	11	12	3	3	1,286	1		1,287	5.57	6.48
(%)	(0.8)	(0.8)	(0.6)	(0.5)	(0.4)	(0.5)	(0.3)	(0.6)	(0.3)	(0.9)	(0.6)	(0.4)		(0.6)		
妊娠腎/妊娠中毒症	1	6	1	1		1	2				12			12	10.00	10.46
(%)	(0.0)	(0.0)	(0.0)	(0.0)		(0.0)	(0.1)				(0.0)			(0.0)		
その他の分類不能の腎炎	154	165	177	112	64	69	45	15	17	3	821			821	9.88	9.41
(%)	(0.3)	(0.3)	(0.4)	(0.5)	(0.5)	(1.0)	(1.3)	(0.7)	(1.5)	(0.9)	(0.4)			(0.4)		
多発性嚢胞腎	1,044	1,415	1,598	1,075	561	288	132	42	21	6	6,182	1		6,183	8.06	7.11
(%)	(2.2)	(2.7)	(3.2)	(4.4)	(4.5)	(4.4)	(3.7)	(2.1)	(1.9)	(1.8)	(3.1)	(0.4)		(3.1)		
腎硬化症	6,494	6,190	4,637	1,626	535	158	61	27	10	1	19,739	6		19,745	4.32	4.64
(%)	(13.8)	(11.7)	(9.2)	(6.6)	(4.3)	(2.4)	(1.7)	(1.3)	(0.9)	(0.3)	(9.8)	(2.3)		(9.8)		
悪性高血圧	392	443	480	225	144	52	27	4	3	1	1,771			1,771	6.71	6.37
(%)	(0.8)	(0.8)	(0.9)	(0.9)	(1.2)	(0.8)	(0.8)	(0.2)	(0.3)	(0.3)	(0.9)			(0.9)		
糖尿病性腎症	22,684	26,356	24,061	8,742	2,304	505	100	18	9	1	84,780	27		84,807	4.73	4.35
(%)	(48.3)	(49.7)	(47.6)	(35.7)	(18.5)	(7.7)	(2.8)	(0.9)	(0.8)	(0.3)	(42.2)	(10.5)		(42.1)		
SLE腎炎	174	178	149	96	52	26	13	5	2		695			695	6.87	6.96
(%)	(0.4)	(0.3)	(0.3)	(0.4)	(0.4)	(0.4)	(0.4)	(0.2)	(0.2)		(0.3)			(0.3)		
アミロイド腎	75	62	41	9	7	1	4		1		200			200	4.36	5.82
(%)	(0.2)	(0.1)	(0.1)	(0.0)	(0.1)	(0.0)	(0.1)		(0.1)		(0.1)			(0.1)		
痛風腎	130	182	231	162	119	64	32	10	5	2	937	2		939	9.55	7.88
(%)	(0.3)	(0.3)	(0.5)	(0.7)	(1.0)	(1.0)	(0.9)	(0.5)	(0.4)	(0.6)	(0.5)	(0.8)		(0.5)		
先天性代謝異常による腎不全	21	31	33	28	19	13	9	2	3		159			159	10.52	8.81
(%)	(0.0)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.3)	(0.1)	(0.3)		(0.1)			(0.1)		
腎・尿路結核	15	19	16	18	13	10	8	6		1	106			106	12.08	10.22
(%)	(0.0)	(0.0)	(0.0)	(0.1)	(0.1)	(0.2)	(0.2)	(0.3)		(0.3)	(0.1)			(0.1)		
腎・尿路結石	86	77	103	56	32	16	5	1	3		379			379	7.36	7.01
(%)	(0.2)	(0.1)	(0.2)	(0.2)	(0.3)	(0.2)	(0.1)	(0.0)	(0.3)		(0.2)			(0.2)		
腎・尿路腫瘍	233	213	163	55	22	9	3	2	1		701			701	4.53	5.19
(%)	(0.5)	(0.4)	(0.3)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)		(0.3)			(0.3)		
閉塞性尿路障害	124	113	107	53	39	21	15	9		1	482			482	7.41	7.89
(%)	(0.3)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.4)	(0.4)		(0.3)	(0.2)			(0.2)		
骨髄腫	98	64	14	6	1		1				184			184	2.28	3.28
(%)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)		(0.0)				(0.1)			(0.1)		
腎形成不全	64	68	80	69	51	34	22	17	6		411			411	11.12	9.39
(%)	(0.1)	(0.1)	(0.2)	(0.3)	(0.4)	(0.5)	(0.6)	(0.8)	(0.5)		(0.2)			(0.2)		
不明	5,034	4,974	4,419	2,067	1,020	454	209	122	51	26	18,376	188		18,564	5.98	6.48
(%)	(10.7)	(9.4)	(8.7)	(8.4)	(8.2)	(6.9)	(5.9)	(6.1)	(4.6)	(7.6)	(9.1)	(72.9)		(9.2)		
再導入	236	270	288	188	141	100	108	68	26	9	1,434			1,434	11.20	10.23
(%)	(0.5)	(0.5)	(0.6)	(0.8)	(1.1)	(1.5)	(3.0)	(3.4)	(2.3)	(2.6)	(0.7)			(0.7)		
その他	1,376	1,325	1,150	538	253	137	59	49	25	8	4,920	12		4,932	6.08	6.90
(%)	(2.9)	(2.5)	(2.3)	(2.2)	(2.0)	(2.1)	(1.7)	(2.4)	(2.2)	(2.3)	(2.4)	(4.7)		(2.4)		
合計	46,946	52,988	50,568	24,516	12,426	6,591	3,560	2,013	1,116	341	201,065	258	1	201,324	6.81	7.16
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		
記載なし	6	4	1	1				1			13			13	4.62	8.36
総計	46,952	52,992	50,569	24,517	12,426	6,591	3,560	2,014	1,116	341	201,078	258	1	201,337	6.81	7.16

数値下のかつこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査