

表 221 2015年末患者，透析歴，原疾患別人数(女性)

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	4,432	5,841	7,652	6,304	4,997	3,493	2,201	1,326	725	213	37,184	13		37,197	11.96	9.58
(%)	(19.8)	(22.8)	(28.1)	(40.6)	(54.7)	(64.2)	(69.4)	(69.7)	(74.0)	(77.2)	(33.3)	(9.7)		(33.2)		
慢性腎盂腎炎	228	272	352	249	184	118	89	60	17	4	1,573			1,573	10.75	9.27
(%)	(1.0)	(1.1)	(1.3)	(1.6)	(2.0)	(2.2)	(2.8)	(3.2)	(1.7)	(1.4)	(1.4)			(1.4)		
急速進行性糸球体腎炎	325	317	286	140	60	42	14	3	3	1	1,191			1,191	6.03	6.42
(%)	(1.4)	(1.2)	(1.1)	(0.9)	(0.7)	(0.8)	(0.4)	(0.2)	(0.3)	(0.4)	(1.1)			(1.1)		
妊娠腎/妊娠中毒症	73	117	220	238	260	231	187	132	60	15	1,533			1,533	17.23	10.19
(%)	(0.3)	(0.5)	(0.8)	(1.5)	(2.8)	(4.2)	(5.9)	(6.9)	(6.1)	(5.4)	(1.4)			(1.4)		
その他の分類不能の腎炎	116	125	133	77	44	38	25	18	6	2	584			584	9.27	9.16
(%)	(0.5)	(0.5)	(0.5)	(0.5)	(0.5)	(0.7)	(0.8)	(0.9)	(0.6)	(0.7)	(0.5)			(0.5)		
多発性嚢胞腎	877	1,012	1,286	910	562	261	108	41	12	3	5,072	1		5,073	8.48	7.20
(%)	(3.9)	(3.9)	(4.7)	(5.9)	(6.2)	(4.8)	(3.4)	(2.2)	(1.2)	(1.1)	(4.5)	(0.7)		(4.5)		
腎硬化症	3,136	3,059	2,478	896	305	124	44	10	3	2	10,057	3		10,060	4.62	4.89
(%)	(14.0)	(11.9)	(9.1)	(5.8)	(3.3)	(2.3)	(1.4)	(0.5)	(0.3)	(0.7)	(9.0)	(2.2)		(9.0)		
悪性高血圧	189	206	211	107	67	41	22	1	1	2	847			847	7.19	7.05
(%)	(0.8)	(0.8)	(0.8)	(0.7)	(0.7)	(0.8)	(0.7)	(0.1)	(0.1)	(0.7)	(0.8)			(0.8)		
糖尿病性腎症	8,666	10,299	10,316	4,346	1,358	362	80	22	7	3	35,459	12		35,471	5.28	4.80
(%)	(38.6)	(40.2)	(37.9)	(28.0)	(14.9)	(6.7)	(2.5)	(1.2)	(0.7)	(1.1)	(31.7)	(9.0)		(31.7)		
SLE腎炎	286	296	318	226	174	100	70	46	14	2	1,532			1,532	9.69	8.91
(%)	(1.3)	(1.2)	(1.2)	(1.5)	(1.9)	(1.8)	(2.2)	(2.4)	(1.4)	(0.7)	(1.4)			(1.4)		
アミロイド腎	81	72	63	32	9	5					262			262	4.95	5.03
(%)	(0.4)	(0.3)	(0.2)	(0.2)	(0.1)	(0.1)					(0.2)			(0.2)		
痛風腎	26	24	31	22	14	16	2	1			136			136	8.98	7.66
(%)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.3)	(0.1)	(0.1)			(0.1)			(0.1)		
先天性代謝異常による腎不全	17	21	31	15	12	6	7	4	2		115			115	10.51	9.37
(%)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)		(0.1)			(0.1)		
腎・尿路結核	8	17	17	10	13	11	2	8	3		89			89	13.16	10.49
(%)	(0.0)	(0.1)	(0.1)	(0.1)	(0.1)	(0.2)	(0.1)	(0.4)	(0.3)		(0.1)			(0.1)		
腎・尿路結石	45	49	53	19	16	13	5	1	1		202			202	7.45	7.59
(%)	(0.2)	(0.2)	(0.2)	(0.1)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)		(0.2)			(0.2)		
腎・尿路腫瘍	59	50	58	28	7	6		1			209			209	5.72	5.53
(%)	(0.3)	(0.2)	(0.2)	(0.2)	(0.1)	(0.1)		(0.1)			(0.2)			(0.2)		
閉塞性尿路障害	37	52	55	40	27	18	13	3	2	1	248			248	9.78	8.54
(%)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.3)	(0.4)	(0.2)	(0.2)	(0.4)	(0.2)			(0.2)		
骨髄腫	55	27	19	5				1			107			107	2.80	4.09
(%)	(0.2)	(0.1)	(0.1)	(0.0)				(0.1)			(0.1)			(0.1)		
腎形成不全	35	44	42	36	34	29	13	8	5	3	249			249	11.86	10.03
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.4)	(0.5)	(0.4)	(0.4)	(0.5)	(1.1)	(0.2)			(0.2)		
不明	2,792	2,846	2,670	1,396	746	359	183	149	79	11	11,231	102		11,333	6.89	7.37
(%)	(12.5)	(11.1)	(9.8)	(9.0)	(8.2)	(6.6)	(5.8)	(7.8)	(8.1)	(4.0)	(10.1)	(76.1)		(10.1)		
再導入	143	133	168	107	56	52	39	32	12	7	749			749	10.32	9.98
(%)	(0.6)	(0.5)	(0.6)	(0.7)	(0.6)	(1.0)	(1.2)	(1.7)	(1.2)	(2.5)	(0.7)			(0.7)		
その他	799	764	772	339	184	114	66	36	28	7	3,109	3		3,112	6.90	7.66
(%)	(3.6)	(3.0)	(2.8)	(2.2)	(2.0)	(2.1)	(2.1)	(1.9)	(2.9)	(2.5)	(2.8)	(2.2)		(2.8)		
合計	22,425	25,643	27,231	15,542	9,129	5,439	3,170	1,903	980	276	111,738	134		111,872	8.22	8.14
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)		
記載なし	2	3	1	1		1					8			8	6.50	7.48
総計	22,427	25,646	27,232	15,543	9,129	5,440	3,170	1,903	980	276	111,746	134		111,880	8.22	8.14

数値下のかつこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査