

表 226 2015年末患者、治療方法(新分類)、年齢別

治療方法 (新分類)	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
施設血液透析	3	2	8	51	150	450	1,191	2,746	6,042	9,612	13,567	18,278	29,078	43,982	38,561	35,834	28,945	15,412	4,213	595	248,720	4	1	248,725	68.57	12.32	
(%)	(8.1)	(7.1)	(21.1)	(49.0)	(70.8)	(68.8)	(71.7)	(72.3)	(72.0)	(72.4)	(73.1)	(74.6)	(76.7)	(78.6)	(80.8)	(83.3)	(85.4)	(86.5)	(86.2)	(88.5)	(79.4)	(100.0)	(100.0)	(79.4)			
Off-line HDF				2	1	8	31	79	154	252	374	497	754	1,009	745	657	487	207	64	11	5,332			5,332	66.34	12.22	
(%)				(1.9)	(0.5)	(1.2)	(1.9)	(2.1)	(1.8)	(1.9)	(2.0)	(2.0)	(2.0)	(1.8)	(1.6)	(1.5)	(1.4)	(1.2)	(1.3)	(1.6)	(1.7)			(1.7)			
On-line HDF			1	9	29	153	311	720	1,687	2,580	3,503	4,301	6,083	8,304	6,378	4,973	3,404	1,628	419	44	44,527			44,527	65.13	12.49	
(%)			(2.6)	(8.7)	(13.7)	(23.4)	(18.7)	(19.0)	(20.1)	(19.4)	(18.9)	(17.5)	(16.0)	(14.8)	(13.4)	(11.6)	(10.0)	(9.1)	(8.6)	(6.5)	(14.2)			(14.2)			
Push/Pull HDF					1			2	4	8	6	11	13	24	26	6	5	4			110			110	64.36	11.80	
(%)					(0.5)			(0.1)	(0.0)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.1)	(0.0)	(0.0)	(0.0)			(0.0)			(0.0)			
ハイオフィльтраーション							1	5	5	8	13	16	34	56	43	42	22	16	6		267			267	68.49	11.79	
(%)							(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)		(0.1)			(0.1)			
IHDF				1	5	5	14	25	77	109	170	249	382	620	608	534	410	253	69	9	3,540			3,540	69.50	12.04	
(%)				(1.0)	(2.4)	(0.8)	(0.8)	(0.7)	(0.9)	(0.8)	(0.9)	(1.0)	(1.0)	(1.1)	(1.3)	(1.2)	(1.2)	(1.4)	(1.4)	(1.3)	(1.1)			(1.1)			
血液濾過									1		1		2	2	3	4	3				16			16	69.94	11.87	
(%)									(0.0)		(0.0)		(0.0)	(0.0)	(0.0)	(0.0)	(0.0)				(0.0)			(0.0)			
血液吸着透析 (リクセル等使用)								2	1	19	53	123	267	400	276	165	58	10	2		1,376			1,376	67.20	7.61	
(%)								(0.1)	(0.0)	(0.1)	(0.3)	(0.5)	(0.7)	(0.7)	(0.6)	(0.4)	(0.2)	(0.1)	(0.0)		(0.4)			(0.4)			
在宅血液透析					1	3	9	26	63	96	76	99	87	53	21	12	6	2	1		555			555	54.49	10.80	
(%)					(0.5)	(0.5)	(0.5)	(0.7)	(0.8)	(0.7)	(0.4)	(0.4)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)		(0.2)			(0.2)			
腹膜透析 (CAPD)		1		4	5	22	46	102	177	296	401	472	663	891	657	510	383	190	73	9	4,902			4,902	64.92	13.12	
(%)		(3.6)		(3.8)	(2.4)	(3.4)	(2.8)	(2.7)	(2.1)	(2.2)	(2.2)	(1.9)	(1.7)	(1.6)	(1.4)	(1.2)	(1.1)	(1.1)	(1.5)	(1.3)	(1.6)			(1.6)			
腹膜透析 (APD)	29	20	21	27	12	11	32	56	132	199	272	297	401	458	299	240	151	78	38	2	2,775			2,775	60.88	16.27	
(%)	(78.4)	(71.4)	(55.3)	(26.0)	(5.7)	(1.7)	(1.9)	(1.5)	(1.6)	(1.5)	(1.5)	(1.2)	(1.1)	(0.8)	(0.6)	(0.6)	(0.4)	(0.4)	(0.8)	(0.3)	(0.9)			(0.9)			
腹膜透析 (CCPD)	5	5	8	10	8	2	26	35	54	98	118	167	161	182	111	55	29	13	3	2	1,092			1,092	58.15	14.48	
(%)	(13.5)	(17.9)	(21.1)	(9.6)	(3.8)	(0.3)	(1.6)	(0.9)	(0.6)	(0.7)	(0.6)	(0.7)	(0.4)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.3)	(0.3)			(0.3)			
合計	37	28	38	104	212	654	1,661	3,798	8,397	13,277	18,554	24,510	37,925	55,981	47,728	43,032	33,903	17,813	4,888	672	313,212	4	1	313,217	67.86	12.49	
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		
記載なし																											
総計	37	28	38	104	212	654	1,661	3,798	8,397	13,277	18,554	24,510	37,925	55,981	47,728	43,032	33,903	17,813	4,888	672	313,212	4	1	313,217	67.86	12.49	

数値下のかっこ内は列方向の合計に対する％です。
*腹膜透析 (CCPD) は、APDと手動のバック交換の両方を行うもの
(C)Japanese Society for Dialysis Therapy
2015年調査