

表 241 2015年死亡患者(性別、原疾患)

原疾患	男性	女性	合計	記載なし	総計
慢性糸球体腎炎	4,100	2,534	6,634		6,634
(%)	(21.6)	(25.1)	(22.8)		(22.8)
慢性腎盂腎炎	121	138	259		259
(%)	(0.6)	(1.4)	(0.9)		(0.9)
急速進行性糸球体腎炎	197	147	344		344
(%)	(1.0)	(1.5)	(1.2)		(1.2)
妊娠腎/妊娠中毒症	1	76	77		77
(%)	(0.0)	(0.8)	(0.3)		(0.3)
その他の分類不能の腎炎	48	35	83		83
(%)	(0.3)	(0.3)	(0.3)		(0.3)
多発性嚢胞腎	348	282	630		630
(%)	(1.8)	(2.8)	(2.2)		(2.2)
腎硬化症	2,176	1,096	3,272		3,272
(%)	(11.4)	(10.9)	(11.2)		(11.2)
悪性高血圧	120	58	178		178
(%)	(0.6)	(0.6)	(0.6)		(0.6)
糖尿病性腎症	8,266	3,565	11,831		11,831
(%)	(43.5)	(35.4)	(40.7)		(40.7)
SLE腎炎	82	126	208		208
(%)	(0.4)	(1.2)	(0.7)		(0.7)
アミロイド腎	34	56	90		90
(%)	(0.2)	(0.6)	(0.3)		(0.3)
痛風腎	70	7	77		77
(%)	(0.4)	(0.1)	(0.3)		(0.3)
先天性代謝異常による腎不全	7	5	12		12
(%)	(0.0)	(0.0)	(0.0)		(0.0)
腎・尿路結核	9	7	16		16
(%)	(0.0)	(0.1)	(0.1)		(0.1)
腎・尿路結石	30	23	53		53
(%)	(0.2)	(0.2)	(0.2)		(0.2)
腎・尿路腫瘍	107	36	143		143
(%)	(0.6)	(0.4)	(0.5)		(0.5)
閉塞性尿路障害	38	17	55		55
(%)	(0.2)	(0.2)	(0.2)		(0.2)
骨髄腫	73	33	106		106
(%)	(0.4)	(0.3)	(0.4)		(0.4)
腎形成不全	9	7	16		16
(%)	(0.0)	(0.1)	(0.1)		(0.1)
不明	2,433	1,455	3,888		3,888
(%)	(12.8)	(14.4)	(13.4)		(13.4)
再導入	70	38	108		108
(%)	(0.4)	(0.4)	(0.4)		(0.4)
その他	669	340	1,009		1,009
(%)	(3.5)	(3.4)	(3.5)		(3.5)
合計	19,008	10,081	29,089		29,089
(%)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし					
総計	19,008	10,081	29,089		29,089

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査