

表 244 2015年死亡患者(透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1 年 ～	2 年 ～	5 年 ～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	194	163	217	382	1,145	1,527	1,057		1,187	714	6,586	19	29	6,634
(%)	(13.7)	(14.0)	(14.8)	(15.5)	(16.7)	(19.6)	(27.0)		(45.0)	(66.2)	(22.9)	(15.0)	(18.1)	(22.8)
慢性腎盂腎炎	11	14	10	28	43	47	39		41	26	259			259
(%)	(0.8)	(1.2)	(0.7)	(1.1)	(0.6)	(0.6)	(1.0)		(1.6)	(2.4)	(0.9)			(0.9)
急速進行性糸球体腎炎	50	36	30	30	105	52	24		10	4	341	2	1	344
(%)	(3.5)	(3.1)	(2.1)	(1.2)	(1.5)	(0.7)	(0.6)		(0.4)	(0.4)	(1.2)	(1.6)	(0.6)	(1.2)
妊娠腎/妊娠中毒症	1	1		2	3	13	8		22	27	77			77
(%)	(0.1)	(0.1)		(0.1)	(0.0)	(0.2)	(0.2)		(0.8)	(2.5)	(0.3)			(0.3)
その他の分類不能の腎炎	5	5	9	9	15	15	7		11	7	83			83
(%)	(0.4)	(0.4)	(0.6)	(0.4)	(0.2)	(0.2)	(0.2)		(0.4)	(0.6)	(0.3)			(0.3)
多発性嚢胞腎	16	13	12	27	108	145	128		138	35	622	3	5	630
(%)	(1.1)	(1.1)	(0.8)	(1.1)	(1.6)	(1.9)	(3.3)		(5.2)	(3.2)	(2.2)	(2.4)	(3.1)	(2.2)
腎硬化症	168	151	219	376	995	854	309		162	21	3,255	3	14	3,272
(%)	(11.9)	(13.0)	(15.0)	(15.2)	(14.5)	(11.0)	(7.9)		(6.1)	(1.9)	(11.3)	(2.4)	(8.8)	(11.2)
悪性高血圧	10	4	10	17	48	42	20		15	8	174	1	3	178
(%)	(0.7)	(0.3)	(0.7)	(0.7)	(0.7)	(0.5)	(0.5)		(0.6)	(0.7)	(0.6)	(0.8)	(1.9)	(0.6)
糖尿病性腎症	373	377	527	1,031	3,134	3,813	1,752		677	46	11,730	29	72	11,831
(%)	(26.4)	(32.4)	(36.0)	(41.7)	(45.6)	(49.0)	(44.8)		(25.7)	(4.3)	(40.7)	(22.8)	(45.0)	(40.7)
SLE腎炎	18	21	10	19	30	44	31		23	8	204	2	2	208
(%)	(1.3)	(1.8)	(0.7)	(0.8)	(0.4)	(0.6)	(0.8)		(0.9)	(0.7)	(0.7)	(1.6)	(1.3)	(0.7)
アミロイド腎	15	6	6	11	24	18	3		4	2	89		1	90
(%)	(1.1)	(0.5)	(0.4)	(0.4)	(0.3)	(0.2)	(0.1)		(0.2)	(0.2)	(0.3)		(0.6)	(0.3)
痛風腎	1	1	2	5	9	24	16		15	4	77			77
(%)	(0.1)	(0.1)	(0.1)	(0.2)	(0.1)	(0.3)	(0.4)		(0.6)	(0.4)	(0.3)			(0.3)
先天性代謝異常による腎不全				1	2	4			4	1	12			12
(%)				(0.0)	(0.0)	(0.1)			(0.2)	(0.1)	(0.0)			(0.0)
腎・尿路結核	1				1	5	2		1	6	16			16
(%)	(0.1)				(0.0)	(0.1)	(0.1)		(0.0)	(0.6)	(0.1)			(0.1)
腎・尿路結石	5		3	1	11	14	12		5	2	53			53
(%)	(0.4)		(0.2)	(0.0)	(0.2)	(0.2)	(0.3)		(0.2)	(0.2)	(0.2)			(0.2)
腎・尿路腫瘍	13	14	10	20	34	35	12		3	1	142		1	143
(%)	(0.9)	(1.2)	(0.7)	(0.8)	(0.5)	(0.4)	(0.3)		(0.1)	(0.1)	(0.5)		(0.6)	(0.5)
閉塞性尿路障害	2	9	4	3	8	16	3		8	2	55			55
(%)	(0.1)	(0.8)	(0.3)	(0.1)	(0.1)	(0.2)	(0.1)		(0.3)	(0.2)	(0.2)			(0.2)
骨髄腫	22	23	14	16	21	9	1				106			106
(%)	(1.6)	(2.0)	(1.0)	(0.6)	(0.3)	(0.1)	(0.0)				(0.4)			(0.4)
腎形成不全	1		1	2	5	2	2		2	1	16			16
(%)	(0.1)		(0.1)	(0.1)	(0.1)	(0.0)	(0.1)		(0.1)	(0.1)	(0.1)			(0.1)
不明	333	232	259	380	892	903	401		264	130	3,794	63	31	3,888
(%)	(23.5)	(19.9)	(17.7)	(15.4)	(13.0)	(11.6)	(10.2)		(10.0)	(12.0)	(13.2)	(49.6)	(19.4)	(13.4)
再導入	9	4	11	7	22	19	8		12	15	107	1		108
(%)	(0.6)	(0.3)	(0.8)	(0.3)	(0.3)	(0.2)	(0.2)		(0.5)	(1.4)	(0.4)	(0.8)		(0.4)
その他	167	89	108	103	219	185	79		35	19	1,004	4	1	1,009
(%)	(11.8)	(7.7)	(7.4)	(4.2)	(3.2)	(2.4)	(2.0)		(1.3)	(1.8)	(3.5)	(3.1)	(0.6)	(3.5)
合計	1,415	1,163	1,462	2,470	6,874	7,786	3,914		2,639	1,079	28,802	127	160	29,089
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし														
総計	1,415	1,163	1,462	2,470	6,874	7,786	3,914		2,639	1,079	28,802	127	160	29,089

数値下のかっこ内は列方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査