

表 249 2015年死亡患者(男性 透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1 年 ～	2 年 ～	5 年 ～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	131	107	134	251	737	949	633		717	412	4,071	11	18	4,100
(%)	(14.2)	(13.4)	(13.0)	(14.4)	(15.9)	(18.7)	(26.1)		(45.1)	(68.6)	(21.6)	(13.9)	(17.3)	(21.6)
慢性腎盂腎炎	6	9	4	18	21	24	15		12	12	121			121
(%)	(0.6)	(1.1)	(0.4)	(1.0)	(0.5)	(0.5)	(0.6)		(0.8)	(2.0)	(0.6)			(0.6)
急速進行性糸球体腎炎	33	20	19	17	51	33	15		4	3	195	1	1	197
(%)	(3.6)	(2.5)	(1.8)	(1.0)	(1.1)	(0.7)	(0.6)		(0.3)	(0.5)	(1.0)	(1.3)	(1.0)	(1.0)
妊娠腎/妊娠中毒症		1									1			1
(%)		(0.1)									(0.0)			(0.0)
その他の分類不能の腎炎	5	4	8	5	8	6	3		2	7	48			48
(%)	(0.5)	(0.5)	(0.8)	(0.3)	(0.2)	(0.1)	(0.1)		(0.1)	(1.2)	(0.3)			(0.3)
多発性嚢胞腎	10	7	10	13	55	84	66		73	22	340	3	5	348
(%)	(1.1)	(0.9)	(1.0)	(0.7)	(1.2)	(1.7)	(2.7)		(4.6)	(3.7)	(1.8)	(3.8)	(4.8)	(1.8)
腎硬化症	119	106	165	274	661	542	191		95	11	2,164	2	10	2,176
(%)	(12.9)	(13.2)	(16.0)	(15.7)	(14.3)	(10.7)	(7.9)		(6.0)	(1.8)	(11.5)	(2.5)	(9.6)	(11.4)
悪性高血圧	7	3	6	11	31	27	19		9	5	118	1	1	120
(%)	(0.8)	(0.4)	(0.6)	(0.6)	(0.7)	(0.5)	(0.8)		(0.6)	(0.8)	(0.6)	(1.3)	(1.0)	(0.6)
糖尿病性腎症	247	270	398	769	2,255	2,625	1,169		442	23	8,198	20	48	8,266
(%)	(26.7)	(33.7)	(38.6)	(44.1)	(48.6)	(51.8)	(48.1)		(27.8)	(3.8)	(43.5)	(25.3)	(46.2)	(43.5)
SLE腎炎	6	11	6	12	13	12	13		7	1	81		1	82
(%)	(0.6)	(1.4)	(0.6)	(0.7)	(0.3)	(0.2)	(0.5)		(0.4)	(0.2)	(0.4)		(1.0)	(0.4)
アミロイド腎	8	3	5	4	7	5			1	1	34			34
(%)	(0.9)	(0.4)	(0.5)	(0.2)	(0.2)	(0.1)			(0.1)	(0.2)	(0.2)			(0.2)
痛風腎	1	1	1	5	9	21	14		14	4	70			70
(%)	(0.1)	(0.1)	(0.1)	(0.3)	(0.2)	(0.4)	(0.6)		(0.9)	(0.7)	(0.4)			(0.4)
先天性代謝異常による腎不全				1	1	2			2	1	7			7
(%)				(0.1)	(0.0)	(0.0)			(0.1)	(0.2)	(0.0)			(0.0)
腎・尿路結核						4	2		1	2	9			9
(%)						(0.1)	(0.1)		(0.1)	(0.3)	(0.0)			(0.0)
腎・尿路結石	2		2	1	7	10	6		1	1	30			30
(%)	(0.2)		(0.2)	(0.1)	(0.2)	(0.2)	(0.2)		(0.1)	(0.2)	(0.2)			(0.2)
腎・尿路腫瘍	10	10	8	14	27	26	9		3		107			107
(%)	(1.1)	(1.2)	(0.8)	(0.8)	(0.6)	(0.5)	(0.4)		(0.2)		(0.6)			(0.6)
閉塞性尿路障害	1	7	2	2	8	11	3		2	2	38			38
(%)	(0.1)	(0.9)	(0.2)	(0.1)	(0.2)	(0.2)	(0.1)		(0.1)	(0.3)	(0.2)			(0.2)
骨髄腫	17	14	10	14	11	6	1				73			73
(%)	(1.8)	(1.7)	(1.0)	(0.8)	(0.2)	(0.1)	(0.0)				(0.4)			(0.4)
腎形成不全	1		1	1	2	2			2		9			9
(%)	(0.1)		(0.1)	(0.1)	(0.0)	(0.0)			(0.1)		(0.0)			(0.0)
不明	204	164	166	255	579	542	221		172	71	2,374	40	19	2,433
(%)	(22.1)	(20.5)	(16.1)	(14.6)	(12.5)	(10.7)	(9.1)		(10.8)	(11.8)	(12.6)	(50.6)	(18.3)	(12.8)
再導入	6	3	8	3	14	10	6		9	11	70			70
(%)	(0.6)	(0.4)	(0.8)	(0.2)	(0.3)	(0.2)	(0.2)		(0.6)	(1.8)	(0.4)			(0.4)
その他	110	61	77	75	139	128	43		22	12	667	1	1	669
(%)	(11.9)	(7.6)	(7.5)	(4.3)	(3.0)	(2.5)	(1.8)		(1.4)	(2.0)	(3.5)	(1.3)	(1.0)	(3.5)
合計	924	801	1,030	1,745	4,636	5,069	2,429		1,590	601	18,825	79	104	19,008
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし														
総計	924	801	1,030	1,745	4,636	5,069	2,429		1,590	601	18,825	79	104	19,008

数値下のかっこ内は列方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査