

表 250 2015年死亡患者(女性 透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1 年 ～	2 年 ～	5 年 ～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	63	56	83	131	408	578	424		470	302	2,515	8	11	2,534
(%)	(12.8)	(15.5)	(19.2)	(18.1)	(18.2)	(21.3)	(28.6)		(44.8)	(63.2)	(25.2)	(16.7)	(19.6)	(25.1)
慢性腎盂腎炎	5	5	6	10	22	23	24		29	14	138			138
(%)	(1.0)	(1.4)	(1.4)	(1.4)	(1.0)	(0.8)	(1.6)		(2.8)	(2.9)	(1.4)			(1.4)
急速進行性糸球体腎炎	17	16	11	13	54	19	9		6	1	146	1		147
(%)	(3.5)	(4.4)	(2.5)	(1.8)	(2.4)	(0.7)	(0.6)		(0.6)	(0.2)	(1.5)	(2.1)		(1.5)
妊娠腎/妊娠中毒症	1			2	3	13	8		22	27	76			76
(%)	(0.2)			(0.3)	(0.1)	(0.5)	(0.5)		(2.1)	(5.6)	(0.8)			(0.8)
その他の分類不能の腎炎		1	1	4	7	9	4		9		35			35
(%)		(0.3)	(0.2)	(0.6)	(0.3)	(0.3)	(0.3)		(0.9)		(0.4)			(0.3)
多発性嚢胞腎	6	6	2	14	53	61	62		65	13	282			282
(%)	(1.2)	(1.7)	(0.5)	(1.9)	(2.4)	(2.2)	(4.2)		(6.2)	(2.7)	(2.8)			(2.8)
腎硬化症	49	45	54	102	334	312	118		67	10	1,091	1	4	1,096
(%)	(10.0)	(12.4)	(12.5)	(14.1)	(14.9)	(11.5)	(7.9)		(6.4)	(2.1)	(10.9)	(2.1)	(7.1)	(10.9)
悪性高血圧	3	1	4	6	17	15	1		6	3	56		2	58
(%)	(0.6)	(0.3)	(0.9)	(0.8)	(0.8)	(0.6)	(0.1)		(0.6)	(0.6)	(0.6)		(3.6)	(0.6)
糖尿病性腎症	126	107	129	262	879	1,188	583		235	23	3,532	9	24	3,565
(%)	(25.7)	(29.6)	(29.9)	(36.1)	(39.3)	(43.7)	(39.3)		(22.4)	(4.8)	(35.4)	(18.8)	(42.9)	(35.4)
SLE腎炎	12	10	4	7	17	32	18		16	7	123	2	1	126
(%)	(2.4)	(2.8)	(0.9)	(1.0)	(0.8)	(1.2)	(1.2)		(1.5)	(1.5)	(1.2)	(4.2)	(1.8)	(1.2)
アミロイド腎	7	3	1	7	17	13	3		3	1	55		1	56
(%)	(1.4)	(0.8)	(0.2)	(1.0)	(0.8)	(0.5)	(0.2)		(0.3)	(0.2)	(0.6)		(1.8)	(0.6)
痛風腎			1			3	2		1		7			7
(%)			(0.2)			(0.1)	(0.1)		(0.1)		(0.1)			(0.1)
先天性代謝異常による腎不全					1	2			2		5			5
(%)					(0.0)	(0.1)			(0.2)		(0.1)			(0.0)
腎・尿路結核	1				1	1				4	7			7
(%)	(0.2)				(0.0)	(0.0)				(0.8)	(0.1)			(0.1)
腎・尿路結石	3		1		4	4	6		4	1	23			23
(%)	(0.6)		(0.2)		(0.2)	(0.1)	(0.4)		(0.4)	(0.2)	(0.2)			(0.2)
腎・尿路腫瘍	3	4	2	6	7	9	3			1	35		1	36
(%)	(0.6)	(1.1)	(0.5)	(0.8)	(0.3)	(0.3)	(0.2)			(0.2)	(0.4)		(1.8)	(0.4)
閉塞性尿路障害	1	2	2	1		5			6		17			17
(%)	(0.2)	(0.6)	(0.5)	(0.1)		(0.2)			(0.6)		(0.2)			(0.2)
骨髄腫	5	9	4	2	10	3					33			33
(%)	(1.0)	(2.5)	(0.9)	(0.3)	(0.4)	(0.1)					(0.3)			(0.3)
腎形成不全				1	3		2			1	7			7
(%)				(0.1)	(0.1)		(0.1)			(0.2)	(0.1)			(0.1)
不明	129	68	93	125	313	361	180		92	59	1,420	23	12	1,455
(%)	(26.3)	(18.8)	(21.5)	(17.2)	(14.0)	(13.3)	(12.1)		(8.8)	(12.3)	(14.2)	(47.9)	(21.4)	(14.4)
再導入	3	1	3	4	8	9	2		3	4	37	1		38
(%)	(0.6)	(0.3)	(0.7)	(0.6)	(0.4)	(0.3)	(0.1)		(0.3)	(0.8)	(0.4)	(2.1)		(0.4)
その他	57	28	31	28	80	57	36		13	7	337	3		340
(%)	(11.6)	(7.7)	(7.2)	(3.9)	(3.6)	(2.1)	(2.4)		(1.2)	(1.5)	(3.4)	(6.3)		(3.4)
合計	491	362	432	725	2,238	2,717	1,485		1,049	478	9,977	48	56	10,081
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし														
総計	491	362	432	725	2,238	2,717	1,485		1,049	478	9,977	48	56	10,081

数値下のかっこ内は列方向の合計に対する%です。
 (C)Japanese Society for Dialysis Therapy
 2015年調査