

表 301 2015年死亡患者死亡原因分類（原疾患との関係）

原疾患	心不全	脳血管障害	感染症	消化管出血	悪性腫瘍	悪液質/尿毒症	心筋梗塞	カリウム中毒/頓死	肝硬変症	自殺/拒否	腸閉塞	肺血栓/肺塞栓	災害・事故死	その他	不明	合計	記載なし	総計
慢性糸球体腎炎	1,898	444	1,373	116	629	322	233	157	68	59	95	18	26	605	583	6,626	8	6,634
(%)	(28.6)	(6.7)	(20.7)	(1.8)	(9.5)	(4.9)	(3.5)	(2.4)	(1.0)	(0.9)	(1.4)	(0.3)	(0.4)	(9.1)	(8.8)	(100.0)		
慢性腎盂腎炎	59	16	63	2	42	13	8	3	2		7		1	20	23	259		259
(%)	(22.8)	(6.2)	(24.3)	(0.8)	(16.2)	(5.0)	(3.1)	(1.2)	(0.8)		(2.7)		(0.4)	(7.7)	(8.9)	(100.0)		
急速進行性糸球体腎炎	75	20	117	6	27	21	9	7	4	1	3	1		27	26	344		344
(%)	(21.8)	(5.8)	(34.0)	(1.7)	(7.8)	(6.1)	(2.6)	(2.0)	(1.2)	(0.3)	(0.9)	(0.3)		(7.8)	(7.6)	(100.0)		
妊娠腎/妊娠中毒症	20	11	17	5	2	2	2	5			1		1	5	6	77		77
(%)	(26.0)	(14.3)	(22.1)	(6.5)	(2.6)	(2.6)	(2.6)	(6.5)			(1.3)		(1.3)	(6.5)	(7.8)	(100.0)		
その他の分類不能の腎炎	15	4	24		9	3	2	5				1		8	11	83		83
(%)	(18.1)	(4.8)	(28.9)		(10.8)	(3.8)	(2.4)	(6.0)				(1.2)		(9.6)	(13.3)	(100.0)		
多発性囊胞腎	125	56	164	11	62	23	18	12	12	4	7	3	7	72	54	630		630
(%)	(19.8)	(8.9)	(26.0)	(1.7)	(9.8)	(3.7)	(2.9)	(1.9)	(1.9)	(0.6)	(1.1)	(0.5)	(1.1)	(11.4)	(8.6)	(100.0)		
腎硬化症	793	193	797	61	303	199	119	73	17	28	34	5	10	354	292	3,268	4	3,272
(%)	(24.3)	(5.6)	(24.4)	(1.9)	(9.3)	(6.1)	(3.6)	(2.2)	(0.5)	(0.9)	(1.0)	(0.2)	(0.3)	(10.8)	(8.9)	(100.0)		
悪性高血圧	56	13	27	5	20	9	7	3	2	2	3			14	17	178		178
(%)	(31.5)	(7.3)	(15.2)	(2.8)	(11.2)	(5.1)	(3.9)	(1.7)	(1.1)	(1.1)	(1.7)			(7.9)	(9.6)	(100.0)		
糖尿病性腎症	3,247	798	2,545	144	943	493	680	362	101	79	100	30	62	978	1,262	11,824	7	11,831
(%)	(27.5)	(6.7)	(21.5)	(1.2)	(8.0)	(4.2)	(5.8)	(3.1)	(0.9)	(0.7)	(0.8)	(0.3)	(0.5)	(8.3)	(10.7)	(100.0)		
SLC腎炎	56	16	49	3	18	6	6	4	3		2	1		26	18	208		208
(%)	(26.9)	(7.7)	(23.8)	(1.4)	(8.7)	(2.9)	(2.9)	(1.9)	(1.4)		(1.0)	(0.5)		(12.5)	(8.7)	(100.0)		
アミロイド腎	23	3	22	2	13	5	2	2				2		11	5	90		90
(%)	(25.6)	(3.3)	(24.4)	(2.2)	(14.4)	(5.6)	(2.2)	(2.2)			(2.2)			(12.2)	(5.6)	(100.0)		
痛風腎	23	3	18	2	8	2	3	1		1				8	8	77		77
(%)	(29.9)	(3.9)	(23.4)	(2.6)	(10.4)	(2.6)	(3.9)	(1.3)		(1.3)				(10.4)	(10.4)	(100.0)		
先天性代謝異常による腎不全	4		4		1		2							1		12		12
(%)	(33.3)		(33.3)		(8.3)		(16.7)							(8.3)		(100.0)		
腎・尿路結核	4		2		1	4				1				1	3	16		16
(%)	(25.0)		(12.5)		(6.3)	(25.0)				(6.3)				(6.3)	(18.8)	(100.0)		
腎・尿路結石	10	8	13	1	3	3	3				1		1	4	6	53		53
(%)	(18.9)	(15.1)	(24.5)	(1.9)	(5.7)	(5.7)	(5.7)				(1.9)		(1.9)	(7.5)	(11.3)	(100.0)		
腎・尿路腫瘍	18	9	24		65	5	4	1	1			1		8	7	143		143
(%)	(12.6)	(6.3)	(16.8)		(45.5)	(3.5)	(2.8)	(0.7)	(0.7)			(0.7)		(5.6)	(4.9)	(100.0)		
閉塞性尿路障害	10	5	15	1	16	1	1	1						4	1	55		55
(%)	(18.2)	(9.1)	(27.3)	(1.8)	(29.1)	(1.8)	(1.8)	(1.8)						(7.3)	(1.8)	(100.0)		
骨髄腫	14		15	1	63	1	1							6	5	106		106
(%)	(13.2)		(14.2)	(0.9)	(59.4)	(0.9)	(0.9)							(5.7)	(4.7)	(100.0)		
腎形成不全	2	1	4		4						2			2	1	16		16
(%)	(12.5)	(6.3)	(25.0)		(25.0)						(12.5)			(12.5)	(6.3)	(100.0)		
不明	880	247	864	51	345	177	116	89	45	28	41	8	16	458	517	3,882	6	3,888
(%)	(22.7)	(6.4)	(22.3)	(1.3)	(8.9)	(4.6)	(3.0)	(2.3)	(1.2)	(0.7)	(1.1)	(0.2)	(0.4)	(11.8)	(13.3)	(100.0)		
再導入	24	13	30		9	2	4	4	2	2	2	1	1	9	5	108		108
(%)	(22.2)	(12.0)	(27.8)		(8.3)	(1.9)	(3.7)	(3.7)	(1.9)	(1.9)	(1.9)	(0.9)	(0.9)	(8.3)	(4.6)	(100.0)		
その他	188	66	206	10	134	48	25	19	23	7	7	3	2	153	118	1,009		1,009
(%)	(18.6)	(6.5)	(20.4)	(1.0)	(13.3)	(4.8)	(2.5)	(1.9)	(2.3)	(0.7)	(0.7)	(0.3)	(0.2)	(15.2)	(11.7)	(100.0)		
合計	7,544	1,916	6,393	421	2,717	1,339	1,245	748	280	212	307	72	128	2,774	2,968	29,064	25	29,089
(%)	(26.0)	(6.6)	(22.0)	(1.4)	(9.3)	(4.6)	(4.3)	(2.6)	(1.0)	(0.7)	(1.1)	(0.2)	(0.4)	(9.5)	(10.2)	(100.0)		
記載なし																		
(%)																		
総計	7,544	1,916	6,393	421	2,717	1,339	1,245	748	280	212	307	72	128	2,774	2,968	29,064	25	29,089
(%)	(26.0)	(6.6)	(22.0)	(1.4)	(9.3)	(4.6)	(4.3)	(2.6)	(1.0)	(0.7)	(1.1)	(0.2)	(0.4)	(9.5)	(10.2)	(100.0)		

数値下のかっこ内は行方向の合計に対する％です。
©Japanese Society for Dialysis Therapy
2015年調査