

表 317 2015年死亡患者死亡原因分類(年齢別, 原疾患別)原疾患＝腎・尿路腫瘍

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)													1 (20.0)			2 (5.9)	4 (12.5)	8 (25.8)	1 (16.7)	2 (66.7)	18 (12.6)			18 (12.6)	85.33	8.16
脳血管障害 (%)															2 (11.8)	4 (11.8)	1 (3.1)	2 (6.5)			9 (6.3)			9 (6.3)	79.33	5.43
感染症 (%)															3 (17.6)	5 (14.7)	9 (28.1)	3 (9.7)	3 (50.0)	1 (33.3)	24 (16.8)			24 (16.8)	82.21	6.14
消化管出血 (%)																										
悪性腫瘍 (%)												1 (50.0)	3 (60.0)	10 (76.9)	10 (58.8)	18 (52.9)	11 (34.4)	11 (35.5)	1 (16.7)		65 (45.5)			65 (45.5)	76.74	7.71
悪液質/尿毒症 (%)														1 (7.7)		1 (2.9)	1 (3.1)	2 (6.5)			5 (3.5)			5 (3.5)	80.00	8.06
心筋梗塞 (%)												1 (50.0)				1 (2.9)	1 (3.1)	1 (3.2)			4 (2.8)			4 (2.8)	75.75	12.39
カリウム中毒/頓死 (%)																			1 (16.7)		1 (0.7)			1 (0.7)	91.00	
肝硬変症 (%)																1 (2.9)					1 (0.7)			1 (0.7)	78.00	
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																		1 (3.2)			1 (0.7)			1 (0.7)	85.00	
災害・事故死 (%)																										
その他 (%)													1 (20.0)		1 (5.9)	1 (2.9)	4 (12.5)	1 (3.2)			8 (5.6)			8 (5.6)	77.63	7.17
不明 (%)														2 (15.4)	1 (5.9)	1 (2.9)	1 (3.1)	2 (6.5)			7 (4.9)			7 (4.9)	76.86	7.47
合計 (%)												2 (100.0)	5 (100.0)	13 (100.0)	17 (100.0)	34 (100.0)	32 (100.0)	31 (100.0)	6 (100.0)	3 (100.0)	143 (100.0)			143 (100.0)	79.21	7.98
記載なし																										
総計												2	5	13	17	34	32	31	6	3	143			143	79.21	7.98

数値下のかっこ内は列方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査