

表 318 2015年死亡患者死亡原因分類(年齢別、原疾患別)原疾患=閉塞性尿路障害

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)							1 (100.0)							1 (20.0)	2 (25.0)	1 (11.1)		3 (25.0)	2 (66.7)		10 (18.2)			10 (18.2)	76.20	17.68	
脳血管障害 (%)														1 (12.5)	3 (33.3)			1 (33.3)		5 (9.1)			5 (9.1)	79.60	6.73		
感染症 (%)														1 (20.0)	1 (12.5)	3 (33.3)	4 (33.3)	5 (41.7)			15 (27.3)			15 (27.3)	79.60	8.57	
消化管出血 (%)																1 (8.3)				1 (1.8)			1 (1.8)	83.00			
悪性腫瘍 (%)														3 (60.0)	4 (50.0)	2 (22.2)	4 (33.3)	1 (8.3)			16 (29.1)			16 (29.1)	72.75	11.62	
悪液質/尿毒症 (%)																				1 (1.8)			1 (1.8)	83.00			
心筋梗塞 (%)																				1 (1.8)			1 (1.8)	42.00			
カリウム中毒/頓死 (%)																				1 (1.8)			1 (1.8)	64.00			
肝硬変症 (%)																											
自殺/拒否 (%)																											
腸閉塞 (%)																											
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)																				2 (16.7)	2 (16.7)		4 (7.3)		4 (7.3)	84.50	3.87
不明 (%)																				1 (8.3)		1 (1.8)		1 (1.8)	85.00		
合計 (%)							1 (100.0)		1 (100.0)	2 (100.0)		1 (100.0)	5 (100.0)	8 (100.0)	9 (100.0)	12 (100.0)	12 (100.0)	3 (100.0)		55 (100.0)			55 (100.0)	76.60	12.38		
記載なし																											
総計							1		1	2		1	1	5	8	9	12	12	3		55			55	76.60	12.38	

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査