

表 324 2015年死亡患者死亡原因分類(治療方法別)

死因	施設血液透析	血液透析濾過	血液濾過	血液吸着透析	在宅血液透析	腹膜透析	離脱	生体腎移植	献腎移植	合計	記載なし	総計
心不全	4,234	496	2	10	2	88				4,832	2,712	7,544
(%)	(26.1)	(29.5)	(25.0)	(19.2)	(40.0)	(22.7)				(26.3)	(25.4)	(26.0)
脳血管障害	1,057	119	1	4	1	31				1,213	703	1,916
(%)	(6.5)	(7.1)	(12.5)	(7.7)	(20.0)	(8.0)				(6.6)	(6.6)	(6.6)
感染症	3,516	319		13	1	105				3,954	2,439	6,393
(%)	(21.6)	(19.0)		(25.0)	(20.0)	(27.1)				(21.5)	(22.8)	(22.0)
消化管出血	235	25		2		6				268	153	421
(%)	(1.4)	(1.5)		(3.8)		(1.5)				(1.5)	(1.4)	(1.4)
悪性腫瘍	1,462	108	2	4	1	19				1,596	1,121	2,717
(%)	(9.0)	(6.4)	(25.0)	(7.7)	(20.0)	(4.9)				(8.7)	(10.5)	(9.3)
悪液質/尿毒症	739	42				19				800	539	1,339
(%)	(4.5)	(2.5)				(4.9)				(4.4)	(5.0)	(4.6)
心筋梗塞	653	94	2	2		18				769	476	1,245
(%)	(4.0)	(5.6)	(25.0)	(3.8)		(4.6)				(4.2)	(4.5)	(4.3)
カリウム中毒/頓死	402	68		3		16				489	259	748
(%)	(2.5)	(4.0)		(5.8)		(4.1)				(2.7)	(2.4)	(2.6)
肝硬変症	144	11		1		8				164	116	280
(%)	(0.9)	(0.7)		(1.9)		(2.1)				(0.9)	(1.1)	(1.0)
自殺/拒否	110	17		1						128	84	212
(%)	(0.7)	(1.0)		(1.9)						(0.7)	(0.8)	(0.7)
腸閉塞	162	27		2		7				198	109	307
(%)	(1.0)	(1.6)		(3.8)		(1.8)				(1.1)	(1.0)	(1.1)
肺血栓/肺塞栓	42	5				1				48	24	72
(%)	(0.3)	(0.3)				(0.3)				(0.3)	(0.2)	(0.2)
災害・事故死	73	10		1		1				85	43	128
(%)	(0.4)	(0.6)		(1.9)		(0.3)				(0.5)	(0.4)	(0.4)
その他	1,628	138		5		35				1,806	968	2,774
(%)	(10.0)	(8.2)		(9.6)		(9.0)				(9.8)	(9.1)	(9.5)
不明	1,792	201	1	4		34				2,032	936	2,968
(%)	(11.0)	(12.0)	(12.5)	(7.7)		(8.8)				(11.1)	(8.8)	(10.2)
合計	16,249	1,680	8	52	5	388				18,382	10,682	29,064
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)				(100.0)	(100.0)	(100.0)
記載なし	6	15								21	4	25
総計	16,255	1,695	8	52	5	388				18,403	10,686	29,089

数値下のカッコ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査