

表 328 2015年死亡患者死亡原因の詳細(原疾患との関係)

死因	慢性 糸球体腎炎	慢性 腎盂腎炎	急速進行 性糸球体腎炎	妊娠腎/妊娠 中毒症	その他の分類 不能の腎炎	多発性嚢胞腎	腎硬化症	悪性高血圧	糖尿病性腎症	SLE腎炎	アミロイド腎	痛風腎
心不全	1,579	45	55	15	12	96	635	41	2,703	40	15	17
(%)	(25.5)	(0.7)	(0.9)	(0.2)	(0.2)	(1.6)	(10.3)	(0.7)	(43.7)	(0.6)	(0.2)	(0.3)
肺水腫(溢水)	78	2	2	1	1	3	26	7	130	3	1	1
(%)	(25.2)	(0.6)	(0.6)	(0.3)	(0.3)	(1.0)	(8.4)	(2.3)	(42.1)	(1.0)	(0.3)	(0.3)
急性心筋梗塞(発症30日以内死亡)	172	7	7	2	2	13	82	7	537	6	2	1
(%)	(18.0)	(0.7)	(0.7)	(0.2)	(0.2)	(1.4)	(8.6)	(0.7)	(56.1)	(0.6)	(0.2)	(0.1)
虚血性心疾患(急性心筋梗塞以外)	61	1	2			5	37		143			2
(%)	(21.3)	(0.3)	(0.7)			(1.7)	(12.9)		(49.8)			(0.7)
不整脈、伝導障害	77	8	3	1		8	43	4	192	6	3	3
(%)	(19.1)	(2.0)	(0.7)	(0.2)		(2.0)	(10.7)	(1.0)	(47.6)	(1.5)	(0.7)	(0.7)
心内膜炎および弁膜症	58	2	5		1	6	22		62	2	2	2
(%)	(30.4)	(1.0)	(2.6)		(0.5)	(3.1)	(11.5)		(32.5)	(1.0)	(1.0)	(1.0)
その他の心疾患	106	2	10	3	1	12	67	4	160	5	2	
(%)	(23.4)	(0.4)	(2.2)	(0.7)	(0.2)	(2.6)	(14.8)	(0.9)	(35.3)	(1.1)	(0.4)	
くも膜下出血	31	1	3			13	13		44	1		
(%)	(22.3)	(0.7)	(2.2)			(9.4)	(9.4)		(31.7)	(0.7)		
脳内出血	206	7	7	6	2	22	61	6	394	9		1
(%)	(24.0)	(0.8)	(0.8)	(0.7)	(0.2)	(2.6)	(7.1)	(0.7)	(45.8)	(1.0)		(0.1)
脳梗塞	161	7	8	4	2	15	82	6	298	5	3	2
(%)	(22.1)	(1.0)	(1.1)	(0.5)	(0.3)	(2.1)	(11.3)	(0.8)	(40.9)	(0.7)	(0.4)	(0.3)
その他の脳血管疾患	46	1	2	1		6	27	1	62	1		
(%)	(24.3)	(0.5)	(1.1)	(0.5)		(3.2)	(14.3)	(0.5)	(32.8)	(0.5)		
敗血症	502	28	45	7	7	80	229	10	1,046	23	14	4
(%)	(21.1)	(1.2)	(1.9)	(0.3)	(0.3)	(3.4)	(9.6)	(0.4)	(44.0)	(1.0)	(0.6)	(0.2)
中枢神経系感染症	10	1	1			1	7	1	13	1		
(%)	(27.0)	(2.7)	(2.7)			(2.7)	(18.9)	(2.7)	(35.1)	(2.7)		
肺炎	720	26	66	6	13	52	481	14	1,242	21	4	12
(%)	(21.7)	(0.8)	(2.0)	(0.2)	(0.4)	(1.6)	(14.5)	(0.4)	(37.5)	(0.6)	(0.1)	(0.4)
インフルエンザ	10	1	1	1		2	9		15			
(%)	(23.8)	(2.4)	(2.4)	(2.4)		(4.8)	(21.4)		(35.7)			
尿路感染症	5					3	2		13			1
(%)	(17.2)					(10.3)	(6.9)		(44.8)			(3.4)
消化管・胆道系感染症・腹膜炎	62	4	1	2	2	9	31	1	75	3	2	
(%)	(25.3)	(1.6)	(0.4)	(0.8)	(0.8)	(3.7)	(12.7)	(0.4)	(30.6)	(1.2)	(0.8)	
劇症(急性)ウイルス肝炎	2					1			3			
(%)	(33.3)					(16.7)			(50.0)			
結核	9		1			1	6		16			1
(%)	(16.7)		(1.9)			(1.9)	(11.1)		(29.6)			(1.9)
ヒト免疫不全ウイルス[HIV]感染症	1											
(%)	(100.0)											
その他の感染症	52	3	2	1	2	15	32	1	122	1	2	
(%)	(17.9)	(1.0)	(0.7)	(0.3)	(0.7)	(5.2)	(11.0)	(0.3)	(42.1)	(0.3)	(0.7)	
中枢神経系の悪性新生物	2						1	1	10			
(%)	(8.3)						(4.2)	(4.2)	(41.7)			
呼吸器系の悪性新生物	113	7	10		1	12	74	4	189	2	1	4
(%)	(21.9)	(1.4)	(1.9)		(0.2)	(2.3)	(14.4)	(0.8)	(36.7)	(0.4)	(0.2)	(0.8)
肝癌	89	4	1		2	8	47	3	196	3	1	1
(%)	(20.9)	(0.9)	(0.2)		(0.5)	(1.9)	(11.0)	(0.7)	(46.0)	(0.7)	(0.2)	(0.2)
肝癌以外の消化器系の悪性新生物	159	10	8	1	3	15	81	4	260	5	1	1
(%)	(24.0)	(1.5)	(1.2)	(0.2)	(0.5)	(2.3)	(12.2)	(0.6)	(39.2)	(0.8)	(0.2)	(0.2)
乳房の悪性新生物	13	4	1				2		14			
(%)	(31.0)	(9.5)	(2.4)				(4.8)		(33.3)			
性器の悪性新生物	27	2				2	12	1	24	1		1
(%)	(26.5)	(2.0)				(2.0)	(11.8)	(1.0)	(23.5)	(1.0)		(1.0)

表 328 2015年死亡患者死亡原因の詳細(原疾患との関係)

死因	慢性 糸球体腎炎	慢性 腎盂腎炎	急速進行 性糸球体腎炎	妊娠腎/妊娠 中毒症	その他の分類 不能の腎炎	多発性嚢胞腎	腎硬化症	悪性高血圧	糖尿病性腎症	SLE腎炎	アミロイド腎	痛風腎
腎の悪性新生物	85	4				10	17	2	31	3	2	
(%)	(39.0)	(1.8)				(4.6)	(7.8)	(0.9)	(14.2)	(1.4)	(0.9)	
内分泌腺の悪性新生物	17		1			1	5		13		1	
(%)	(34.7)		(2.0)			(2.0)	(10.2)		(26.5)		(2.0)	
造血・リンパ組織の悪性新生物	37	3	2	1		7	17	1	52	2	7	1
(%)	(14.9)	(1.2)	(0.8)	(0.4)		(2.8)	(6.8)	(0.4)	(20.9)	(0.8)	(2.8)	(0.4)
その他の悪性新生物	87	8	4		3	7	47	4	154	2		
(%)	(20.3)	(1.9)	(0.9)		(0.7)	(1.6)	(11.0)	(0.9)	(35.9)	(0.5)		
ウイルス性肝硬変	21	1				3	5	1	44			
(%)	(21.0)	(1.0)				(3.0)	(5.0)	(1.0)	(44.0)			
ウイルス性以外の肝硬変	47	1	4			9	12	1	57	3		
(%)	(26.1)	(0.6)	(2.2)			(5.0)	(6.7)	(0.6)	(31.7)	(1.7)		
腸の血行障害	55	5	3			4	16	2	65	1	1	
(%)	(29.7)	(2.7)	(1.6)			(2.2)	(8.6)	(1.1)	(35.1)	(0.5)	(0.5)	
イレウス	36	2		1		3	17	1	35	1		
(%)	(32.4)	(1.8)		(0.9)		(2.7)	(15.3)	(0.9)	(31.5)	(0.9)		
消化管出血	116	2	6	5		11	61	5	144	3	2	2
(%)	(27.6)	(0.5)	(1.4)	(1.2)		(2.6)	(14.5)	(1.2)	(34.2)	(0.7)	(0.5)	(0.5)
被嚢性腹膜硬化症	4						1				1	
(%)	(36.4)						(9.1)				(9.1)	
その他の消化器疾患	35		2		2	14	27	1	46	2	1	1
(%)	(19.7)		(1.1)		(1.1)	(7.9)	(15.2)	(0.6)	(25.8)	(1.1)	(0.6)	(0.6)
肺梗塞、肺塞栓症	18		1		1	3	5		30	1		
(%)	(25.0)		(1.4)		(1.4)	(4.2)	(6.9)		(41.7)	(1.4)		
悪液質	181	8	9	1	2	11	115	6	237	3	1	1
(%)	(25.9)	(1.1)	(1.3)	(0.1)	(0.3)	(1.6)	(16.4)	(0.9)	(33.9)	(0.4)	(0.1)	(0.1)
尿毒症	91	4	7	1		7	46	2	151	2	3	
(%)	(23.2)	(1.0)	(1.8)	(0.3)		(1.8)	(11.7)	(0.5)	(38.5)	(0.5)	(0.8)	
認知症	18		2			1	13	1	24	1		
(%)	(27.3)		(3.0)			(1.5)	(19.7)	(1.5)	(36.4)	(1.5)		
その他の悪液質/尿毒症	32	1	3		1	4	25		81		1	1
(%)	(17.7)	(0.6)	(1.7)		(0.6)	(2.2)	(13.8)		(44.8)		(0.6)	(0.6)
高カリウム血症	21	2	1	1	3	1	6		52			1
(%)	(18.9)	(1.8)	(0.9)	(0.9)	(2.7)	(0.9)	(5.4)		(46.8)			(0.9)
原因不明の突然死	136	1	6	4	2	11	67	3	310	4	2	
(%)	(21.4)	(0.2)	(0.9)	(0.6)	(0.3)	(1.7)	(10.5)	(0.5)	(48.7)	(0.6)	(0.3)	
自殺	31						11	2	32			1
(%)	(35.2)						(12.5)	(2.3)	(36.4)			(1.1)
治療拒否(透析拒否)	28		1			4	17		47			
(%)	(22.6)		(0.8)			(3.2)	(13.7)		(37.9)			
災害・事故死	26	1		1	1	7	10		62			
(%)	(20.3)	(0.8)		(0.8)	(0.8)	(5.5)	(7.8)		(48.4)			
その他	570	20	25	5	6	58	327	13	932	24	10	7
(%)	(22.0)	(0.8)	(1.0)	(0.2)	(0.2)	(2.2)	(12.6)	(0.5)	(35.9)	(0.9)	(0.4)	(0.3)
合計	6,043	236	318	71	72	576	2,976	161	10,562	190	85	69
(%)	(23.2)	(0.9)	(1.2)	(0.3)	(0.3)	(2.2)	(11.4)	(0.6)	(40.5)	(0.7)	(0.3)	(0.3)
記載なし	591	23	26	6	11	54	296	17	1,269	18	5	8
総計	6,634	259	344	77	83	630	3,272	178	11,831	208	90	77

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査

表 328 2015年死亡患者死亡原因の詳細(原疾患との関係)

死因	先天性代謝異常に基づく腎不全	腎・尿路結核	腎・尿路結石	腎・尿路腫瘍	閉塞性尿路障害	骨髄腫	腎形成不全	不明	再導入	その他	合計	記載なし	総計
心不全	3	4	8	12	7	10	2	714	21	154	6,188		6,188
(%)	(0.0)	(0.1)	(0.1)	(0.2)	(0.1)	(0.2)	(0.0)	(11.5)	(0.3)	(2.5)	(100.0)		
肺水腫(溢水)				1	2	1		41		9	309		309
(%)				(0.3)	(0.6)	(0.3)		(13.3)		(2.9)	(100.0)		
急性心筋梗塞(発症30日以内死亡)	1		2	3	1			89	4	20	958		958
(%)	(0.1)		(0.2)	(0.3)	(0.1)			(9.3)	(0.4)	(2.1)	(100.0)		
虚血性心疾患(急性心筋梗塞以外)	1		1	1		1		27		5	287		287
(%)	(0.3)		(0.3)	(0.3)		(0.3)		(9.4)		(1.7)	(100.0)		
不整脈、伝導障害			1	2		2		41	1	8	403		403
(%)			(0.2)	(0.5)		(0.5)		(10.2)	(0.2)	(2.0)	(100.0)		
心内膜炎および弁膜症				1				24		4	191		191
(%)				(0.5)				(12.6)		(2.1)	(100.0)		
その他の心疾患	1		1	2	1	1		60	2	13	453		453
(%)	(0.2)		(0.2)	(0.4)	(0.2)	(0.2)		(13.2)	(0.4)	(2.9)	(100.0)		
くも膜下出血				2	1			25	1	4	139		139
(%)				(1.4)	(0.7)			(18.0)	(0.7)	(2.9)	(100.0)		
脳内出血			3	4	1			98	4	29	860		860
(%)			(0.3)	(0.5)	(0.1)			(11.4)	(0.5)	(3.4)	(100.0)		
脳梗塞			4	1	2			95	7	26	728		728
(%)			(0.5)	(0.1)	(0.3)			(13.0)	(1.0)	(3.6)	(100.0)		
その他の脳血管疾患			1	2	1		1	29	1	7	189		189
(%)			(0.5)	(1.1)	(0.5)		(0.5)	(15.3)	(0.5)	(3.7)	(100.0)		
敗血症	2	2	5	5	5	7	1	262	12	80	2,376		2,376
(%)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)	(0.3)	(0.0)	(11.0)	(0.5)	(3.4)	(100.0)		
中枢神経系感染症								1		1	37		37
(%)								(2.7)		(2.7)	(100.0)		
肺炎	2		5	14	9	5	2	501	15	103	3,313		3,313
(%)	(0.1)		(0.2)	(0.4)	(0.3)	(0.2)	(0.1)	(15.1)	(0.5)	(3.1)	(100.0)		
インフルエンザ								3			42		42
(%)								(7.1)			(100.0)		
尿路感染症					1	1		1		2	29		29
(%)					(3.4)	(3.4)		(3.4)		(6.9)	(100.0)		
消化管・胆道系感染症・腹膜炎			1	4		1	1	35	2	9	245		245
(%)			(0.4)	(1.6)		(0.4)	(0.4)	(14.3)	(0.8)	(3.7)	(100.0)		
劇症(急性)ウイルス肝炎											6		6
(%)											(100.0)		
結核								18		2	54		54
(%)								(33.3)		(3.7)	(100.0)		
ヒト免疫不全ウイルス[HIV]感染症											1		1
(%)											(100.0)		
その他の感染症			2	1		1		43	1	9	290		290
(%)			(0.7)	(0.3)		(0.3)		(14.8)	(0.3)	(3.1)	(100.0)		
中枢神経系の悪性新生物	1			1				7		1	24		24
(%)	(4.2)			(4.2)				(29.2)		(4.2)	(100.0)		
呼吸器系の悪性新生物				8	1		2	59		28	515		515
(%)				(1.6)	(0.2)		(0.4)	(11.5)		(5.4)	(100.0)		
肝癌						1	1	50		19	426		426
(%)						(0.2)	(0.2)	(11.7)		(4.5)	(100.0)		
肝癌以外の消化器系の悪性新生物		1	1	1	5	1		79	2	25	663		663
(%)		(0.2)	(0.2)	(0.2)	(0.8)	(0.2)		(11.9)	(0.3)	(3.8)	(100.0)		
乳房の悪性新生物								3		5	42		42
(%)								(7.1)		(11.9)	(100.0)		
性器の悪性新生物				4	4		1	16	2	5	102		102
(%)				(3.9)	(3.9)		(1.0)	(15.7)	(2.0)	(4.9)	(100.0)		

表 328 2015年死亡患者死亡原因の詳細(原疾患との関係)

死因	先天性代謝異常に基づく腎不全	腎・尿路結核	腎・尿路結石	腎・尿路腫瘍	閉塞性尿路障害	骨髄腫	腎形成不全	不明	再導入	その他	合計	記載なし	総計
腎の悪性新生物 (%)				27 (12.4)	1 (0.5)			23 (10.6)	1 (0.5)	12 (5.5)	218 (100.0)		218
内分泌腺の悪性新生物 (%)								8 (16.3)		3 (6.1)	49 (100.0)		49
造血・リンパ組織の悪性新生物 (%)			1 (0.4)	2 (0.8)		53 (21.3)		39 (15.7)	3 (1.2)	21 (8.4)	249 (100.0)		249
その他の悪性新生物 (%)			1 (0.2)	22 (5.1)	5 (1.2)	8 (1.9)		61 (14.2)	1 (0.2)	15 (3.5)	429 (100.0)		429
ウイルス性肝硬変 (%)								19 (19.0)	1 (1.0)	5 (5.0)	100 (100.0)		100
ウイルス性以外の肝硬変 (%)				1 (0.6)				26 (14.4)	1 (0.6)	18 (10.0)	180 (100.0)		180
膵の血行障害 (%)			1 (0.5)					26 (14.1)	2 (1.1)	4 (2.2)	185 (100.0)		185
イレウス (%)								14 (12.6)		1 (0.9)	111 (100.0)		111
消化管出血 (%)			1 (0.2)		1 (0.2)	1 (0.2)		51 (12.1)		10 (2.4)	421 (100.0)		421
被囊性腹膜硬化症 (%)							2 (18.2)	1 (9.1)		2 (18.2)	11 (100.0)		11
その他の消化器疾患 (%)		1 (0.6)		1 (0.6)				35 (19.7)	2 (1.1)	8 (4.5)	178 (100.0)		178
肺梗塞、肺塞栓症 (%)				1 (1.4)				8 (11.1)	1 (1.4)	3 (4.2)	72 (100.0)		72
悪液質 (%)		3 (0.4)	3 (0.4)	2 (0.3)				102 (14.6)	1 (0.1)	14 (2.0)	700 (100.0)		700
尿毒症 (%)				3 (0.8)	1 (0.3)	1 (0.3)		49 (12.5)		24 (6.1)	392 (100.0)		392
認知症 (%)								4 (6.1)		2 (3.0)	66 (100.0)		66
その他の悪液質/尿毒症 (%)		1 (0.6)						22 (12.2)	1 (0.6)	8 (4.4)	181 (100.0)		181
高カリウム血症 (%)					1 (0.9)			20 (18.0)	1 (0.9)	1 (0.9)	111 (100.0)		111
原因不明の突然死 (%)				1 (0.2)				69 (10.8)	3 (0.5)	18 (2.8)	637 (100.0)		637
自殺 (%)								9 (10.2)		2 (2.3)	88 (100.0)		88
治療拒否(透析拒否) (%)		1 (0.8)						19 (15.3)	2 (1.6)	5 (4.0)	124 (100.0)		124
災害・事故死 (%)			1 (0.8)					16 (12.5)	1 (0.8)	2 (1.6)	128 (100.0)		128
その他 (%)	1 (0.0)		4 (0.2)	7 (0.3)	4 (0.2)	6 (0.2)	2 (0.1)	423 (16.3)	7 (0.3)	145 (5.6)	2,596 (100.0)		2,596
合計 (%)	12 (0.0)	13 (0.0)	47 (0.2)	136 (0.5)	54 (0.2)	101 (0.4)	15 (0.1)	3,365 (12.9)	103 (0.4)	891 (3.4)	26,096 (100.0)		26,096
記載なし		3	6	7	1	5	1	523	5	118	2,993		2,993
総計	12	16	53	143	55	106	16	3,888	108	1,009	29,089		29,089

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査