

表 330 1983年以降導入2015年末患者原疾患別人数, 平均年齢

原疾患	人 数	記 載 な し	総 計	平 均	標 準 偏 差
慢性糸球体腎炎	90,257		90,257	66.95	12.60
(%)	(29.2)		(29.2)		
慢性腎盂腎炎	2,846		2,846	65.68	13.97
(%)	(0.9)		(0.9)		
急速進行性糸球体腎炎	2,462		2,462	69.29	12.93
(%)	(0.8)		(0.8)		
妊娠腎/妊娠中毒症	1,424		1,424	64.76	10.50
(%)	(0.5)		(0.5)		
その他分類不能の腎炎	1,366		1,366	61.39	17.03
(%)	(0.4)		(0.4)		
多発性嚢胞腎	11,186		11,186	65.15	11.39
(%)	(3.6)		(3.6)		
腎硬化症	29,779		29,779	74.25	11.74
(%)	(9.6)		(9.6)		
悪性高血圧	2,610		2,610	64.03	14.90
(%)	(0.8)		(0.8)		
糖尿病性腎症	120,249	2	120,251	67.52	11.33
(%)	(38.9)	(50.0)	(38.9)		
SLE腎炎	2,194		2,194	61.65	13.73
(%)	(0.7)		(0.7)		
アミロイド腎	461		461	68.16	11.35
(%)	(0.1)		(0.1)		
痛風腎	1,065		1,065	67.76	11.42
(%)	(0.3)		(0.3)		
先天性代謝異常による腎不全	266		266	49.45	17.42
(%)	(0.1)		(0.1)		
腎・尿路結核	183		183	73.46	8.79
(%)	(0.1)		(0.1)		
腎・尿路結石	576		576	70.88	11.17
(%)	(0.2)		(0.2)		
腎・尿路腫瘍	909		909	72.38	10.54
(%)	(0.3)		(0.3)		
閉塞性尿路障害	723		723	64.13	16.40
(%)	(0.2)		(0.2)		
骨髄腫	291		291	70.76	10.83
(%)	(0.1)		(0.1)		
腎形成不全	639		639	45.13	19.54
(%)	(0.2)		(0.2)		
不明	29,636	2	29,638	70.02	13.15
(%)	(9.6)	(50.0)	(9.6)		
移植後再導入	2,100		2,100	56.65	12.69
(%)	(0.7)		(0.7)		
その他	7,953		7,953	66.22	15.50
(%)	(2.6)		(2.6)		
合計	309,175	4	309,179	67.89	12.54
(%)	(100.0)	(100.0)	(100.0)		
記載なし	21		21	70.52	11.36
総計	309,196	4	309,200	67.89	12.54

数値下のかっこ内は列方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査